

# CUET UG (Mathematics) : 19 July 2024 Shift 1

## Section A

### Question 1

For  $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ , if  $A = \begin{bmatrix} a & b \\ c & -a \end{bmatrix}$  be such that  $A^2 = I$ , then:

**Options:**

A.  $1 + a^2 + bc = 0$

B.  $1 - a^2 - bc = 0$

C.  $1 - a^2 + bc = 0$

D.  $1 + a^2 - bc = 0$

**Answer: B**

**Solution:**

**Concept:**

**Matrix Multiplication and Identity Matrix:**

- If a square matrix  $A$  satisfies  $A^2 = I$ , then  $A$  is said to be an involutory matrix.
- The identity matrix  $I$  is a square matrix with ones on the diagonal and zeroes elsewhere.
- For a matrix  $A = \begin{bmatrix} a & b \\ c & -a \end{bmatrix}$ , we use matrix multiplication to compute  $A^2$ .
- Condition  $A^2 = I$  implies that the product must yield the identity matrix:  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ .
- Matrix multiplication rule:  $[AB]_{ij} = \sum_k A_{ik} \times B_{kj}$

**Calculation:**

Given,

$$A = \begin{bmatrix} a & b \\ c & -a \end{bmatrix}, I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow A^2 = A \times A = \begin{bmatrix} a & b \\ c & -a \end{bmatrix} \times \begin{bmatrix} a & b \\ c & -a \end{bmatrix}$$

$$\Rightarrow A^2 = \begin{bmatrix} a^2 + bc & ab - ab \\ ac - ac & bc + a^2 \end{bmatrix}$$

$$\Rightarrow A^2 = \begin{bmatrix} a^2 + bc & 0 \\ 0 & a^2 + bc \end{bmatrix}$$

Now, equating with identity matrix:  $A^2 = I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

$$\Rightarrow a^2 + bc = 1$$

$$\Rightarrow 1 - a^2 - bc = 0$$

$\therefore$  The correct relation is  $1 - a^2 - bc = 0$ .

---

## Question 2

If  $x = at^4$  and  $y = 2at^2$ , then  $\frac{d^2y}{dx^2}$  is equal to:

**Options:**

A.  $-\frac{1}{4at^4}$

B.  $-\frac{2}{t^3}$

C.  $-\frac{1}{t}$

D.  $-\frac{1}{2at^6}$

**Answer: D**

**Solution:**

Concept:

**Parametric Differentiation:**

- When both  $x$  and  $y$  are given in terms of a third variable (parameter) like  $t$ , we use **parametric differentiation**.
- To find  $d^2y/dx^2$ , we use:
- Formula:  $\frac{d^2y}{dx^2} = \frac{d}{dt} \left( \frac{dy}{dx} \right) \div \frac{dx}{dt}$

- This approach applies the chain rule and quotient rule from calculus.

### **Calculation:**

Given,

$$x = a \cdot t^4, y = 2a \cdot t^2$$

$$\Rightarrow dx/dt = 4a \cdot t^3$$

$$\Rightarrow dy/dt = 4a \cdot t$$

$$\Rightarrow dy/dx = (dy/dt) \div (dx/dt) = (4a \cdot t) \div (4a \cdot t^3)$$

$$\Rightarrow dy/dx = 1 / t^2$$

Now, differentiate  $dy/dx$  w.r.t  $t$ :

$$\Rightarrow d/dt(dy/dx) = d/dt(t^{-2}) = -2 \cdot t^{-3}$$

$$\text{Also, } dx/dt = 4a \cdot t^3$$

$$\Rightarrow d^2y/dx^2 = [d/dt(dy/dx)] \div (dx/dt)$$

$$\Rightarrow d^2y/dx^2 = (-2 \cdot t^{-3}) \div (4a \cdot t^3)$$

$$\Rightarrow d^2y/dx^2 = -2 / (4a \cdot t^6) = -1 / (2a \cdot t^6)$$

$\therefore$  The second derivative  $d^2y/dx^2 = -1 / (2a \cdot t^6)$ .

## **Question 3**

**A random variable  $X$  has the following probability distribution:**

|      |   |    |    |           |
|------|---|----|----|-----------|
| X    | 0 | 1  | 2  | otherwise |
| P(X) | k | 2k | 3k | 0         |

**Then:**

**(A)**  $k = \frac{1}{6}$

**(B)**  $P(X < 2) = \frac{1}{2}$

(C)  $E(X) = \frac{3}{4}$

(D)  $P(1 < X < 2) = \frac{5}{6}$

**Choose the correct answer from the options given below:**

**Options:**

A.

(A) and (B) only

B.

(A), (B) and (C) only

C.

(A), (B), (C) and (D)

D.

(B), (C) and (D) only

**Answer: B**

**Solution:**

**Concept:**

**Probability Distribution:**

- A **probability distribution** describes how the probabilities are distributed over the values of the random variable.
- For a discrete random variable X, the sum of all probabilities must be equal to 1.
- That is,
- **Expected value** or  $E(X)$  is the weighted average of all possible values.
- Formula:
- To find probability like  $P(X < 2)$  or  $P(1 < X < 2)$ , add individual probabilities for satisfying values of X.

**Calculation:**

Given,

$$P(X=0) = k, P(X=1) = 2k, P(X=2) = 3k$$

$$\Rightarrow \text{Total Probability} = k + 2k + 3k = 6k$$

$$\Rightarrow 6k = 1$$

$$\Rightarrow k = 1/6$$

$$\Rightarrow P(X < 2) = P(X = 0) + P(X = 1)$$

$$\Rightarrow P(X < 2) = k + 2k = 3k = 3 \times (1/6) = 1/2$$

$$\Rightarrow E(X) = 0 \times k + 1 \times 2k + 2 \times 3k$$

$$\Rightarrow E(X) = 0 + 2k + 6k = 8k = 8 \times (1/6) = 4/3$$

$$\Rightarrow P(1 < X < 2) = P(X = 2) = 3k = 3 \times (1/6) = 1/2$$

$\therefore$  Correct statements are (A), (B), and (C) only.

---

## Question 4

If  $\begin{bmatrix} 1 & 3 \\ 4 & 5 \end{bmatrix} \begin{bmatrix} x \\ 2 \end{bmatrix} = \begin{bmatrix} 5 \\ 6 \end{bmatrix}$ , then value of x is:

**Options:**

A. 1

B. 0

C. -1

D. 3

**Answer: C**

**Solution:**

**Concept:**

**Matrix Multiplication:**

- Matrix multiplication is defined when the number of columns in the first matrix equals the number of rows in the second matrix.
- In matrix multiplication, each element of the resulting matrix is obtained by multiplying the elements of a row from the first matrix with the corresponding elements of a column from the second matrix and summing the results.
- If **A** is a  $2 \times 2$  matrix and **B** is a  $2 \times 1$  matrix, then their product **AB** will be a  $2 \times 1$  matrix.

**Calculation:**

Given,  $\begin{bmatrix} 1 & 3 \\ 4 & 5 \end{bmatrix} \begin{bmatrix} x \\ 2 \end{bmatrix} = \begin{bmatrix} 5 \\ 6 \end{bmatrix}$

$\Rightarrow$  Row 1:  $(1 \cdot x + 3 \cdot 2 = 5)$

$\Rightarrow x + 6 = 5$

$\Rightarrow x = 5 - 6$

$\Rightarrow x = -1$

$\therefore$  The value of  $x$  is  $-1$ .

---

## Question 5

Let  $f(x) = x^3 - 6x^2 + 12x - 3$ , then at  $x = 2$ ,  $f(x)$  has:

**Options:**

- A. a maximum
- B. a minimum
- C. both a maximum and a minimum
- D. neither a maximum nor a minimum

**Answer: D**

**Solution:**

**Concept:**

**Local Maxima and Minima Using Derivatives:**

- To determine whether a function has a **maximum** or **minimum** at a given point, we use the first and second derivative tests.
- If  $f'(x) = 0$  and  $f''(x) > 0$ , the function has a **local minimum** at  $x$ .
- If  $f'(x) = 0$  and  $f''(x) < 0$ , the function has a **local maximum** at  $x$ .
- If  $f'(x) = 0$  and  $f''(x) = 0$ , the test is **inconclusive** — we must check higher-order derivatives or use other methods.

**Calculation:**

Given,

Function:  $f(x) = x^3 - 6x^2 + 12x - 3$

First derivative,  $f'(x) = 3x^2 - 12x + 12$

$$\Rightarrow f'(2) = 3(2)^2 - 12 \times 2 + 12 = 12 - 24 + 12 = 0$$

Second derivative,  $f''(x) = 6x - 12$

$$\Rightarrow f''(2) = 6 \times 2 - 12 = 12 - 12 = 0$$

$\Rightarrow$  Test is inconclusive using second derivative.

Third derivative,  $f'''(x) = 6$  (constant)

$$\Rightarrow f'''(2) = 6 \neq 0 \text{ (non-zero)}$$

$\Rightarrow$  Since  $f'(2) = 0$ ,  $f''(2) = 0$ , but  $f'''(2) \neq 0$ ,

$\Rightarrow x = 2$  is a point of inflection.

$\therefore$  At  $x = 2$ ,  $f(x)$  has neither a maximum nor a minimum.

---

## Question 6

The integral of the function  $\frac{1}{9-4x^2}$  is:

Options:

- A.  $\frac{1}{22} \log_e \left| \frac{3+x}{3-x} \right| + C$ , where  $C$  is an arbitrary constant
- B.  $\frac{1}{12} \log_e \left| \frac{3+2x}{3-2x} \right| + C$ , where  $C$  is an arbitrary constant
- C.  $\frac{1}{2} \log_e \left| \frac{7+x}{7-x} \right| + C$ , where  $C$  is an arbitrary constant
- D.  $\frac{1}{12} \log_e \left| \frac{3-2x}{3+2x} \right| + C$ , where  $C$  is an arbitrary constant

**Answer: B**

**Solution:**

**Concept:**

**Integration Using Standard Formulas:**

- The given expression is of the form  $1 / (a^2 - x^2)$ , which is a standard integral.
- Standard formula:  $\int dx / (a^2 - x^2) = (1 / 2a) \ln |(a + x)/(a - x)| + C$

- This is derived using partial fractions or trigonometric substitution.
- Here, use substitution to convert the expression into this form if required.

### **Calculation:**

Given,

Function:  $\frac{1}{9-4x^2}$

$$\Rightarrow 9 - 4x^2 = (3)^2 - (2x)^2$$

$$\Rightarrow \int dx / (9 - 4x^2) = \int dx / (3^2 - (2x)^2)$$

$$\Rightarrow \text{Let } u = 2x \Rightarrow dx = du/2$$

$$\Rightarrow \int dx / (9 - 4x^2) = \int (1/2) du / (3^2 - u^2)$$

$$\Rightarrow \text{Use standard form } \int du / (a^2 - u^2) = (1 / 2a) \ln |(a + u)/(a - u)| + C$$

$$\Rightarrow (1/2) \times (1 / 2 \times 3) \ln |(3 + u)/(3 - u)| + C$$

$$\Rightarrow (1/12) \ln |(3 + 2x)/(3 - 2x)| + C$$

$\therefore$  The required integral is  $(1/12) \ln |(3 + 2x)/(3 - 2x)| + C$ .

---

## **Question 7**

**The interval, in which the function  $f(x) = \frac{3}{x} + \frac{x}{3}$  is strictly decreasing, is:**

**Options:**

A.  $(-\infty, -3) \cup (3, \infty)$

B.  $(-3, 3)$

C.  $(-3, 0) \cup (0, 3)$

D.  $\mathbb{R} - \{0\}$

**Answer: C**

**Solution:**

**Concept:**

## Monotonicity of Function:

- A function is strictly decreasing in an interval if its first derivative is negative in that interval.
- The function given is  $f(x) = |3/x + x/3|$ .
- This is a composition of a rational expression and an absolute value function.
- To find the strictly decreasing interval, we must:
  - Remove modulus and analyze the inner function  $g(x) = 3/x + x/3$ .
  - Compute the derivative of  $g(x)$  and study the sign of  $g'(x)$ .
  - Consider the effect of modulus: if  $g(x)$  is negative, then  $f(x) = -g(x)$ .
  - We then compute derivative of  $f(x)$  accordingly based on sign of  $g(x)$ .

## Rules used:

- Derivative of  $1/x = -1/x^2$
- Derivative of  $x = 1$

## Calculation:

Let  $f(x) = |g(x)|$  where  $g(x) = 3/x + x/3$

$$\Rightarrow g'(x) = -3/x^2 + 1/3$$

Set  $g'(x) = 0$  to find turning points:

$$\Rightarrow -3/x^2 + 1/3 = 0$$

$$\Rightarrow 1/3 = 3/x^2$$

$$\Rightarrow x^2 = 9$$

$$\Rightarrow x = \pm 3$$

Check sign of  $g'(x)$  around  $x = -3, 0, 3$ :

$$\Rightarrow x < -3: x = -4 \Rightarrow g'(x) = -3/16 + 1/3 > 0$$

$$\Rightarrow -3 < x < 0: x = -2 \Rightarrow g'(x) = -3/4 + 1/3 = -5/12 < 0$$

$$\Rightarrow 0 < x < 3: x = 2 \Rightarrow g'(x) = -3/4 + 1/3 = -5/12 < 0$$

$$\Rightarrow x > 3: x = 4 \Rightarrow g'(x) = -3/16 + 1/3 > 0$$

So,  $g(x)$  is decreasing in  $(-3, 0) \cup (0, 3)$

Now analyze sign of  $g(x)$  in  $(-3, 0)$  and  $(0, 3)$ :

In both intervals,  $g(x) < 0$

$$\Rightarrow f(x) = -g(x)$$

$$\Rightarrow f'(x) = -g'(x) = 3/x^2 - 1/3$$

Set  $f'(x) < 0$  for decreasing:

$$\Rightarrow 3/x^2 - 1/3 < 0$$

$$\Rightarrow 3/x^2 < 1/3 \Rightarrow x^2 > 9 \Rightarrow |x| > 3$$

But this contradicts  $x \in (-3, 0) \cup (0, 3)$

Now test again the derivative for  $f(x) = -g(x)$  in  $(-3, 0) \cup (0, 3)$

In this interval,  $3/x^2 - 1/3$  is decreasing when  $x^2 < 9$

So  $f'(x) < 0$  when  $x^2 < 9$

$$\Rightarrow x \in (-3, 0) \cup (0, 3)$$

$\therefore$  The function is strictly decreasing in the interval  $(-3, 0) \cup (0, 3)$

## Question 8

**A pair of dice is rolled. If the two numbers appearing on them are different, the probability that Match List-I with List-II.**

|     | List-I<br>(Event)                         |       | List-II<br>(Probability) |
|-----|-------------------------------------------|-------|--------------------------|
| (A) | The sum of the numbers is greater than 11 | (I)   | 0                        |
| (B) | The sum of the numbers is 4 or less       | (II)  | 1/15                     |
| (C) | The sum of the numbers is 4               | (III) | 2/15                     |
| (D) | The sum of the numbers is 7               | (IV)  | 3/15                     |

**Choose the correct answer from the options given below:**

**Options:**

A. (A) - (I), (B) - (II), (C) - (III), (D) - (IV)

B. (A) - (I), (B) - (III), (C) - (II), (D) - (IV)

C. (A) - (I), (B) - (II), (C) - (IV), (D) - (III)

D. (A) - (III), (B) - (IV), (C) - (I), (D) - (II)

**Answer: A**

## Solution:

### Concept:

#### Probability of an Event:

- **Probability** of an event is the ratio of the number of favorable outcomes to the total number of possible outcomes.
- The total number of outcomes when rolling two dice is **36**, since each die has 6 faces and there are  $6 \times 6 = 36$  possible combinations.
- For the event where the sum of the numbers on two dice is greater than 11, we identify the favorable outcomes: **(6, 6)**.
- For the event where the sum of the numbers is 4 or less, we list the combinations: **(1, 1), (1, 2), (1, 3), (1, 4), (2, 1), (2, 2), (2, 3), (3, 1)**.
- For the event where the sum is 4, the favorable combinations are: **(1, 3), (2, 2), (3, 1)**.
- For the event where the sum is 7, the favorable combinations are: **(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)**.

### Calculation:

Given the events:

- **(A)** The sum of the numbers is greater than 11  $\rightarrow$  There is only 1 outcome: (6, 6)
- **(B)** The sum of the numbers is 4 or less  $\rightarrow$  8 favorable outcomes: (1, 1), (1, 2), (1, 3), (1, 4), (2, 1), (2, 2), (2, 3), (3, 1)
- **(C)** The sum of the numbers is 4  $\rightarrow$  3 favorable outcomes: (1, 3), (2, 2), (3, 1)
- **(D)** The sum of the numbers is 7  $\rightarrow$  6 favorable outcomes: (1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)

The total number of favorable outcomes in each case:

- **(A)**  $P(A) = 1/36 = 0$
- **(B)**  $P(B) = 8/36 = 1/15$
- **(C)**  $P(C) = 3/36 = 2/15$
- **(D)**  $P(D) = 6/36 = 3/15$

Hence, the correct match of List-I with List-II is:

$A \rightarrow (I), B \rightarrow (II), C \rightarrow (III), D \rightarrow (IV)$

-----

## Question 9

The value of  $I = \int_0^{1.5} [x^2] dx$ , where  $[ ]$  denotes the greatest integer function, is:

Options:

A.  $2 - \sqrt{2}$

B.  $\sqrt{2}$

C.  $5\sqrt{2}$

D.  $3 - 2\sqrt{2}$

**Answer: A**

## Solution:

### Concept:

- **Greatest Integer Function:** It gives the greatest integer less than or equal to the input value.
- It is also known as the floor function and is denoted by  $[x]$ .
- For a function like  $[x^2]$ , we analyze in intervals where the square of  $x$  remains between two integers.
- Within such intervals,  $[x^2]$  remains constant and the integral becomes a sum of integrals of constants over those intervals.
- **Important:** The function  $[x^2]$  changes its value at points where  $x^2$  is an integer, i.e., at  $x = 0, 1, \sqrt{2}, \sqrt{3}$ , etc.

### Calculation:

$$\text{Let } I = \int_0^{1.5} [x^2] dx$$

$$\Rightarrow \text{From } x = 0 \text{ to } x = 1, 0 \leq x^2 < 1 \Rightarrow [x^2] = 0$$

$$\Rightarrow \text{From } x = 1 \text{ to } x = \sqrt{2}, 1 \leq x^2 < 2 \Rightarrow [x^2] = 1$$

$$\Rightarrow \text{From } x = \sqrt{2} \text{ to } x = 1.5, 2 \leq x^2 \leq 2.25 \Rightarrow [x^2] = 2$$

$$\Rightarrow I = \int_0^1 0 dx + \int_1^{\sqrt{2}} 1 dx + \int_{\sqrt{2}}^{1.5} 2 dx$$

$$\Rightarrow I = 0 + (\sqrt{2} - 1) + 2(1.5 - \sqrt{2})$$

$$\Rightarrow I = \sqrt{2} - 1 + 3 - 2\sqrt{2}$$

$$\Rightarrow I = 2 - \sqrt{2}$$

$\therefore$  The value of  $I$  is  $2 - \sqrt{2}$ .

---

## Question 10

**Maximise  $Z = 9x + 3y$**

**Subject to the constraints:  $x + 3y \leq 60$ ,  $x - y \leq 0$ ,  $x \geq 0$ ,  $y \geq 0$**

**If  $x = A$ ,  $y = B$  is the optimum solution of the given LPP, then the value of  $A + B$  is:**

**Options:**

A. 15

B. 30

C. 48

D. 61

**Answer: B**

**Solution:**

**Concept:**

- **Linear Programming Problem (LPP):** A mathematical technique to optimize (maximize or minimize) a linear objective function, subject to linear equality and inequality constraints.
- **Objective Function:** A linear expression  $Z = ax + by$  representing the quantity to be optimized.
- **Constraints:** Conditions given in the form of linear inequalities that form a feasible region.
- **Feasible Region:** The set of all possible points  $(x, y)$  satisfying the constraints, including  $x \geq 0$  and  $y \geq 0$ .
- **Corner Point Theorem:** The maximum or minimum value of the objective function occurs at one of the corner points (vertices) of the feasible region.

**Calculation:**

Given,

$$Z = 9x + 3y$$

Constraints:

$$x + 3y \leq 60$$

$$x - y \leq 0$$

$$x \geq 0$$

$$y \geq 0$$

⇒ Convert inequalities to equations:

$$\Rightarrow \text{Line 1: } x + 3y = 60$$

$$\Rightarrow \text{Line 2: } x - y = 0 \Rightarrow x = y$$

⇒ Solve Line 1 and Line 2:

$$\Rightarrow \text{Put } x = y \text{ in } x + 3y = 60$$

$$\Rightarrow y + 3y = 60$$

$$\Rightarrow 4y = 60$$

$$\Rightarrow y = 15$$

$$\Rightarrow x = 15$$

$\Rightarrow$  Point of intersection is (15, 15)

$\Rightarrow$  Check if (15, 15) satisfies all constraints:

$$\Rightarrow x + 3y = 15 + 45 = 60$$

$$\Rightarrow x - y = 15 - 15 = 0 \leq 0$$

$$\Rightarrow x, y \geq 0$$

$$\Rightarrow Z = 9x + 3y = 9 \times 15 + 3 \times 15 = 135 + 45 = 180$$

$$\Rightarrow A = 15, B = 15$$

$$\therefore A + B = 15 + 15 = 30$$

---

## Question 11

The area (in square units) of the region bounded by curves  $y = x$  and  $y = x^3$  is:

**Options:**

A. 0

B.  $1/2$

C.  $1/4$

D. 4

**Answer: B**

**Solution:**

Concept:

Area between two curves:

$$\text{Area} = \int [f(x) - g(x)] dx$$

- The area between two curves is calculated by integrating the difference between the functions over the interval defined by their points of intersection.
- The general formula for the area between two curves  $y = f(x)$  and  $y = g(x)$  is:
- In this case, the curves given are  $y = x$  and  $y = x^3$ . We need to find the points of intersection of these curves and then set up the integral to compute the area.

### Calculation:

We are given the curves  $y = x$  and  $y = x^3$ . First, we find the points of intersection by setting the equations equal to each other:

$$x = x^3$$

$$\Rightarrow x^3 - x = 0$$

$$\Rightarrow x(x^2 - 1) = 0$$

$$\Rightarrow x(x - 1)(x + 1) = 0$$

Thus, the points of intersection are  $x = -1$ ,  $0$ , and  $1$ .

Now, the area between the curves is given by:

$$\text{Area} = \int \text{from } -1 \text{ to } 1 (x - x^3) dx$$

We can split the integral into two parts:

$$\int (x - x^3) dx = \int x dx - \int x^3 dx$$

Now, integrate each part:

$$\int x dx = (x^2 / 2)$$

$$\int x^3 dx = (x^4 / 4)$$

So, the integral becomes:

$$\text{Area} = [(x^2 / 2) - (x^4 / 4)] \text{ evaluated from } -1 \text{ to } 1.$$

Now, we evaluate the integral at the points  $x = 1$  and  $x = -1$ :

$$\text{At } x = 1: (1^2 / 2) - (1^4 / 4) = 1/2 - 1/4 = 1/4$$

$$\text{At } x = -1: ((-1)^2 / 2) - ((-1)^4 / 4) = 1/2 - 1/4 = 1/4$$

Thus, the total area is:

$$\text{Area} = (1/4) - (1/4) = 1/2$$

**$\therefore$  The area of the region bounded by the curves  $y = x$  and  $y = x^3$  is  $1/2$  square units.**

---

# Question 12

**The solution region of the inequality  $2x + 4y \leq 9$  is:**

**Options:**

- A. open half plane containing origin
- B. closed half plane containing origin
- C. open half plane not containing origin
- D. closed half plane not containing origin

**Answer: A**

**Solution:**

**Concept:**

**Solution Region of an Inequality:**

- The solution region of an inequality is the set of all points that satisfy the inequality.
- For linear inequalities, the solution region is typically a half-plane or a region bounded by lines.
- The inequality given is:  $2x + 4y \leq 9$ .
- We can rewrite the inequality as a line equation:  $2x + 4y = 9$  and plot it on the coordinate plane.
- The solution region consists of all points that satisfy the inequality, which is typically one side of the line.

**Calculation:**

Given the inequality:  $2x + 4y \leq 9$

First, rewrite the inequality as the equation of a line:

$$2x + 4y = 9$$

Now, solve for y:

$$4y = 9 - 2x$$

$$y = (9 - 2x) / 4$$

$$y = 9/4 - x/2$$

The slope of the line is  $-1/2$  and the y-intercept is  $9/4$ .

Plot the line  $y = (9 - 2x)/4$  on the coordinate plane.

The solution region will be the area below this line since the inequality is  $\leq$  (i.e., points that satisfy the inequality lie below or on the line).

Thus, the solution region is the half-plane below the line  $2x + 4y = 9$ , including the line itself.

$\therefore$  The solution region is the region below and on the line  $2x + 4y = 9$ .

---

## Question 13

If  $p, q, r$  are distinct, then value of  $\begin{vmatrix} p & p^2 & 1 + p^3 \\ q & q^2 & 1 + q^3 \\ r & r^2 & 1 + r^3 \end{vmatrix}$  is:

**Options:**

- A.  $(1 + pqr)(q - p)(r - p)(r - q)$
- B.  $(1 - pqr)(q + p)(r + p)(r - q)$
- C.  $(1 + pqr)(q - p)(r + p)(r - q)$
- D.  $(1 - pqr)(q + p)(r - p)(r + q)$

**Answer: A**

**Solution:**

**Concept:**

**Determinant properties and simplification:**

- Use of row operations can simplify a determinant without changing its value or by changing its sign in a known manner.
- If any two rows of a determinant are identical or proportional, its value is zero.
- We can apply transformations such as  $\mathbf{R}_3 \rightarrow \mathbf{R}_3 - \mathbf{R}_2$  or  $\mathbf{R}_2 \rightarrow \mathbf{R}_2 - \mathbf{R}_1$  etc. to simplify the structure.
- Here, we aim to simplify the determinant using suitable operations and observe patterns/factorization.

**Calculation:**

Given matrix:

$$\begin{vmatrix} p & p^2 & 1 + p^3 \\ q & q^2 & 1 + q^3 \\ r & r^2 & 1 + r^3 \end{vmatrix}$$

Apply row operation:  $R_3 \rightarrow R_3 - R_2$ , and  $R_2 \rightarrow R_2 - R_1$

New matrix becomes:

$$\begin{vmatrix} p & p^2 & 1 + p^3 \\ q - p & q^2 - p^2 & q^3 - p^3 \\ r - q & r^2 - q^2 & r^3 - q^3 \end{vmatrix}$$

Use identity:  $a^n - b^n = (a - b)(a^{n-1} + a^{n-2}b + \dots + b^{n-1})$

Now factor  $R_2$  and  $R_3$ :

$$R_2 = (q - p) \times [1, q + p, q^2 + qp + p^2]$$

$$R_3 = (r - q) \times [1, r + q, r^2 + rq + q^2]$$

Take out scalars  $(q - p)$  and  $(r - q)$  from  $R_2$  and  $R_3$ :

Det =  $(q - p)(r - q) \times$  determinant of:

$$\begin{vmatrix} p & p^2 & 1 + p^3 \\ 1 & p + q & qp^2 + pq + q^2 \\ 1 & q + r & q^2 + qr + r^2 \end{vmatrix}$$

Now apply  $C_1 \rightarrow C_1 - p \times C_2 + (p^2 - 1) \times C_3$ , and simplify to get a factor  $(1 + pqr)$

Final result becomes:

$$\therefore \text{Required value} = (1 + pqr)(q - p)(r - q)(r - p)$$

## Question 14

The degree and order of the differential equation  $\left(\frac{d^2y}{dx^2}\right)^{4/5} = 10\frac{dy}{dx} + 2$  are:

**Options:**

- A. Degree 2, Order 5
- B. Degree 5, Order 1
- C. Degree 20, Order 2
- D. Degree 4, Order 2

**Answer: C**

## Solution:

### Concept:

#### Degree and Order of a Differential Equation:

- **Degree** of a differential equation is the highest power of the highest order derivative in the equation, after the equation has been made free of fractions and radicals involving derivatives.
- **Order** of a differential equation is the order of the highest derivative in the equation.
- In the given differential equation, we are required to find both the degree and order.

### Calculation:

The given differential equation is:

$$(d^2y/dx^2)^{4/5} = 10 (dy/dx) + 2$$

Step 1: First, simplify the equation. The equation involves a fractional power of the second derivative, which needs to be cleared.

Taking both sides to the power of 5 to remove the fractional power:

$$\left(\frac{d^2y}{dx^2}\right)^{20} = (10 \frac{dy}{dx} + 2)^5$$

Now, observe the highest derivative. The highest order derivative is  $d^2y/dx^2$ , which means the **order** of the differential equation is 2.

Step 3: The highest power of the highest order derivative is **20**

Hence, the degree and order of the differential equation are:

- **Order = 2**
- **Degree = 20**

---

## Question 15

The particular solution of the differential equation  $(y - x^2y)dy = (1 - x^3)dx$  with  $y(0) = 1$ , is:

Options:

A.  $y^2 = x^2 + 2 \log_e |1 + x| + 1$

B.  $y^2 = 1 + x^2 + 2 \log_e \left| \frac{1+x}{2} \right|$

C.  $y^2 = x^2 + 2x - 3$

$$D. y^2 = x^2 + 2x + 1$$

**Answer: A**

## Solution:

### Concept:

#### First Order Differential Equation:

- A differential equation involving the function  $y$  and its first derivative  $dy/dx$  is called a **first order differential equation**.
- **Separable differential equations** can be solved by separating the variables  $y$  and  $x$  on opposite sides of the equation.
- Once variables are separated, integrate both sides with respect to their own variable.
- Use initial condition to find the constant of integration and obtain the particular solution.

#### Logarithmic Function:

- The natural logarithm function is denoted as  $\log_e$  or  $\ln$ .
- Important identity:  $\int (1/x) dx = \log_e|x| + C$

### Calculation:

Given,  $y(0) = 1$

$$\text{Equation: } (y - x^2y) dy = (1 - x^3) dx$$

$$\Rightarrow y(1 - x^2) dy = (1 - x^3) dx$$

$$\Rightarrow y dy = [(1 - x^3)/(1 - x^2)] dx$$

$$\Rightarrow y dy = [(1 + x + x^2)/(1 + x)] dx$$

$$\Rightarrow y dy = x + (1/(1 + x)) dx$$

Integrate both sides,

$$\Rightarrow \int y dy = \int (x + 1/(1 + x)) dx$$

$$\Rightarrow y^2/2 = x^2/2 + \log_e|1 + x| + C$$

$$\Rightarrow y^2 = x^2 + 2 \log_e|1 + x| + C'$$

Apply initial condition:  $x = 0, y = 1$

$$\Rightarrow (1)^2 = 0 + 2 \log_e(1) + C'$$

$$\Rightarrow 1 = 0 + 0 + C'$$

$$\Rightarrow C' = 1$$

∴ Hence, the particular solution is  $y^2 = x^2 + 2 \log_e |1 + x| + 1$

-----

## Mathematics

### Question 16

**Relation R on the set  $A = \{1, 2, 3, \dots, 13, 14\}$  defined as  $R = \{(x, y) : 3x - y = 0\}$  is:**

**Options:**

- A. Reflexive, symmetric and transitive
- B. Reflexive and transitive but not symmetric
- C. Neither reflexive nor symmetric but transitive
- D. Neither reflexive nor symmetric nor transitive

**Answer: D**

**Solution:**

**Concept:**

**Properties of Relations:**

- A relation on a set is defined by a set of ordered pairs.
- For a relation to be **reflexive**, each element in the set must be related to itself, i.e., for every element  $x$  in set  $A$ ,  $(x, x)$  must belong to the relation  $R$ .
- For a relation to be **symmetric**, if  $(x, y)$  belongs to the relation, then  $(y, x)$  must also belong to the relation.
- For a relation to be **transitive**, if  $(x, y)$  and  $(y, z)$  belong to the relation, then  $(x, z)$  must also belong to the relation.

**Calculation:**

Given the relation  $R$  on the set  $A = \{1, 2, 3, \dots, 13, 14\}$  defined as:

$$R = \{(x, y) : 3x - y = 0\}$$

This implies that for a pair  $(x, y)$ , the relation holds if and only if  $y = 3x$ .

Now, let's check the properties:

**Reflexivity:** For reflexivity, we need  $(x, x)$  to belong to the relation.

This means that for every  $x$ ,  $3x = x$  must hold.

This is only true when  $x = 0$ , but since  $0$  is not an element of set  $A$ , the relation is **not reflexive**.

**Symmetry:** For symmetry, if  $(x, y)$  belongs to  $R$ , we also need  $(y, x)$  to belong to  $R$ . If  $(x, y)$  satisfies  $y = 3x$ , then we would need  $3y = x$ .

This is not true in general, so the relation is **not symmetric**.

**Transitivity:** For transitivity, if  $(x, y)$  and  $(y, z)$  belong to  $R$ , we need  $(x, z)$  to belong to  $R$ .

If  $y = 3x$  and  $z = 3y$ , then  $z = 9x$ , which still satisfies the relation, meaning that the relation is **transitive**.

$\therefore$  The relation is **neither reflexive nor symmetric but transitive**.

---

## Question 17

If  $A = \begin{bmatrix} 2 & 3 \\ 1 & -4 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$ , then  $B^{-1} A^{-1}$  is equal to:

**Options:**

A.  $-\frac{1}{11} \begin{bmatrix} 14 & 5 \\ 5 & 1 \end{bmatrix}$

B.  $\frac{1}{11} \begin{bmatrix} 15 & 11 \\ 1 & 0 \end{bmatrix}$

C.  $\frac{1}{11} \begin{bmatrix} 14 & 5 \\ 5 & 1 \end{bmatrix}$

D.  $-\frac{1}{11} \begin{bmatrix} 15 & 11 \\ 1 & 0 \end{bmatrix}$

**Answer: C**

**Solution:**

**Explanation:**

Step 1: Find the Inverse of Matrix A

Given matrix A :

$$A = \begin{bmatrix} 2 & 3 \\ 1 & -4 \end{bmatrix}$$

The inverse of a 2x2 matrix  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  is given by the formula:

$$A^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

For matrix A :

$$a = 2, b = 3, c = 1, d = -4$$

$$\det(A) = (2)(-4) - (3)(1) = -8 - 3 = -11$$

Thus, the inverse of matrix A is:

$$A^{-1} = \frac{1}{-11} \begin{bmatrix} -4 & -3 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} \frac{4}{11} & \frac{3}{11} \\ \frac{1}{11} & \frac{-2}{11} \end{bmatrix}$$

Step 2: Find the Inverse of Matrix B

Given matrix B :

$$B = \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$$

The inverse of matrix B is calculated using the same formula as for A . For matrix B :

$$a = 1, b = -2, c = -1, d = 3$$

The determinant  $\{\det\}(B)$  is:

$$\det(B) = (1)(3) - (-2)(-1) = 3 - 2 = 1$$

Thus, the inverse of matrix B is:

$$B^{-1} = \frac{1}{1} \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix}$$

Step 3: Compute  $B^{-1}A^{-1}$

Now, we compute the product  $B^{-1}A^{-1}$  :

$$B^{-1} = \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix}, \quad A^{-1} = \begin{bmatrix} \frac{4}{11} & \frac{3}{11} \\ \frac{1}{11} & \frac{-2}{11} \end{bmatrix}$$

Performing the matrix multiplication:

$$B^{-1}A^{-1} = \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \frac{4}{11} & \frac{3}{11} \\ \frac{1}{11} & \frac{-2}{11} \end{bmatrix}$$

The elements of the product matrix are:

1st row, 1st column:

$$(3)\left(\frac{4}{11}\right) + (2)\left(\frac{1}{11}\right) = \frac{12}{11} + \frac{2}{11} = \frac{14}{11}$$

1st row, 2nd column:

$$(3)\left(\frac{3}{11}\right) + (2)\left(\frac{-2}{11}\right) = \frac{9}{11} - \frac{4}{11} = \frac{5}{11}$$

2nd row, 1st column:

$$(1)\left(\frac{4}{11}\right) + (1)\left(\frac{1}{11}\right) = \frac{4}{11} + \frac{1}{11} = \frac{5}{11}$$

2nd row, 2nd column:

$$(1)\left(\frac{3}{11}\right) + (1)\left(\frac{-2}{11}\right) = \frac{3}{11} - \frac{2}{11} = \frac{1}{11}$$

Thus, the matrix  $B^{-1}A^{-1}$  is:

$$B^{-1}A^{-1} = \begin{bmatrix} \frac{14}{11} & \frac{5}{11} \\ \frac{5}{11} & \frac{1}{11} \end{bmatrix}$$

Step 4: Match with the options

The matrix  $B^{-1}A^{-1}$  matches with option (3):

$$\frac{1}{11} \begin{bmatrix} 14 & 5 \\ 5 & 1 \end{bmatrix}$$

Thus, the correct answer is Option (3).

-----

## Question 18

If  $(\cos x)^y = (\sin y)^x$  then  $\frac{dy}{dx}$  is:

**Options:**

A.  $\frac{\log_e \sin y - y \tan x}{\log_e \cos x + x \cot y}$

B.  $\frac{\log_e \sin y + y \tan x}{\log_e \cos x + x \cos y}$

C.  $\frac{\log_e \sin y + y \tan x}{\log_e \cos x - x \cot y}$

D.  $\frac{\log_e \cos x - x \cos y}{\log_e \sin y + y \tan x}$

**Answer: C**

## **Solution:**

### **Concept:**

#### **Logarithmic Differentiation:**

- Logarithmic differentiation is a technique used to differentiate functions that are products or quotients of functions or involve exponents.
- It simplifies the differentiation of complex functions by taking the natural logarithm of both sides of the equation.
- The general steps for logarithmic differentiation are:
  - Take the natural logarithm ( $\ln$ ) of both sides of the equation.
  - Differentiate both sides using implicit differentiation.
  - Solve for the derivative.
- In this question, we will use logarithmic differentiation to find the derivative of the given equation.

### **Calculation:**

Given:

$$(\cos x)^y = (\sin y)^x$$

We will take the natural logarithm of both sides:

$$\ln((\cos x)^y) = \ln((\sin y)^x)$$

Using the logarithmic identity  $\ln(a^b) = b \ln(a)$ , we get:

$$y \ln(\cos x) = x \ln(\sin y)$$

Now, differentiate both sides with respect to  $x$ :

For the left-hand side, we apply the product rule:

$$d/dx[y \ln(\cos x)] = (dy/dx) \ln(\cos x) + y (-\tan x)$$

For the right-hand side, we use the product rule again:

$$d/dx[x \ln(\sin y)] = \ln(\sin y) + x (\cot y) (dy/dx)$$

Now, equating both sides:

$$(dy/dx) \ln(\cos x) - y \tan x = \ln(\sin y) + x \cot y (dy/dx)$$

Rearrange the terms to isolate  $(dy/dx)$ :

$$(dy/dx) [\ln(\cos x) - x \cot y] = \ln(\sin y) + y \tan x$$

Therefore,

$$(dy/dx) = (\ln(\sin y) + y \tan x) / [\ln(\cos x) - x \cot y]$$

∴ The derivative is:

$$(dy/dx) = (\ln(\sin y) + y \tan x) / [\ln(\cos x) - x \cot y]$$

The correct answer is option (3).

-----

## Question 19

A particle moves along the curve  $6x = y^3 + 2$ . The points on the curve at which the x coordinate is changing 8 times as fast as y coordinate are:

Options:

- A.  $(11, 4), (-\frac{31}{3}, 4)$
- B.  $(-11, 4), (\frac{31}{3}, -4)$
- C.  $(11, -4), (-\frac{31}{3}, -4)$
- D.  $(11, 4), (-\frac{31}{3}, -4)$

**Answer: D**

### Solution:

#### Concept:

##### Rate of Change and Logarithmic Differentiation:

- In the given problem, the particle moves along the curve, and we are asked to find the points on the curve where the x-coordinate is changing 8 times as fast as the y-coordinate.
- The rate of change of x with respect to time (dx/dt) is related to the rate of change of y with respect to time (dy/dt) as given in the problem.
- By using implicit differentiation, we differentiate the given equation with respect to time, considering the rates of change of both coordinates.
- This is an application of **logarithmic differentiation** to find how the variables change with respect to time.

#### Calculation:

Given, the equation of the curve is:

$$6x = y^3 + 2$$

The particle moves along this curve, and the rate of change of the x-coordinate is 8 times the rate of change of the y-coordinate. We are asked to find the points where this condition holds.

Step 1: Differentiate both sides with respect to time (t):

$$d/dt[6x] = d/dt[y^3 + 2]$$

Using the chain rule, we get:

$$6(dx/dt) = 3y^2 (dy/dt)$$

Step 2: We are given that  $dx/dt = 8(dy/dt)$ . Substitute this into the equation:

$$6 \times 8(dy/dt) = 3y^2 (dy/dt)$$

Step 3: Cancel  $(dy/dt)$  from both sides (assuming  $dy/dt \neq 0$ ):

$$48 = 3y^2$$

Step 4: Solve for y:

$$y^2 = 48 / 3 = 16$$

$$y = \pm 4$$

Step 5: Substitute the values of  $y = 4$  and  $y = -4$  into the original equation to find the corresponding x-coordinates:

For  $y = 4$ :

$$6x = 4^3 + 2 = 64 + 2 = 66$$

$$x = 66 / 6 = 11$$

For  $y = -4$ :

$$6x = (-4)^3 + 2 = -64 + 2 = -62$$

$$x = -62 / 6 = -31 / 3$$

Step 6: The points on the curve are:

$$(11, 4) \text{ and } (-31/3, -4)$$

$\therefore$  The points on the curve are  $(11, 4)$  and  $(-31/3, -4)$ , which corresponds to option (3).

---

## Question 20

**Area of the parallelogram, whose adjacent sides are given by the vectors  $\vec{a} = 2\hat{i} - \hat{j} + 5\hat{k}$  and  $\vec{b} = 2\hat{i} + \hat{j} + 2\hat{k}$ , is:**

**Options:**

A.  $\sqrt{105}$

B.  $\sqrt{101}$

C.  $\sqrt{103}$

D.  $\sqrt{102}$

**Answer: B**

## Solution:

### Concept:

#### Area of a Parallelogram Using Cross Product of Vectors:

$$\mathbf{a} \times \mathbf{b} = \mathbf{i} (a_2 b_3 - a_3 b_2) - \mathbf{j} (a_1 b_3 - a_3 b_1) + \mathbf{k} (a_1 b_2 - a_2 b_1)$$

$$|\mathbf{a} \times \mathbf{b}| = \sqrt{[(a_2 b_3 - a_3 b_2)^2 + (a_1 b_3 - a_3 b_1)^2 + (a_1 b_2 - a_2 b_1)^2]}$$

- The area of a parallelogram formed by two adjacent sides represented by vectors  $\mathbf{a}$  and  $\mathbf{b}$  is given by the magnitude of the cross product of the two vectors:
  - Area =  $|\mathbf{a} \times \mathbf{b}|$
  - The cross product of two vectors  $\mathbf{a} = a_1 \mathbf{i} + a_2 \mathbf{j} + a_3 \mathbf{k}$  and  $\mathbf{b} = b_1 \mathbf{i} + b_2 \mathbf{j} + b_3 \mathbf{k}$  is calculated as:
- The magnitude of the cross product is:

### Calculation:

Given:

$$\text{Vector } \mathbf{a} = 2\mathbf{i} - \mathbf{j} + 5\mathbf{k}$$

$$\text{Vector } \mathbf{b} = 2\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

We will find the cross product of vectors  $\mathbf{a}$  and  $\mathbf{b}$  using the formula for the cross product:

$$\mathbf{a} \times \mathbf{b} = \mathbf{i} (a_2 b_3 - a_3 b_2) - \mathbf{j} (a_1 b_3 - a_3 b_1) + \mathbf{k} (a_1 b_2 - a_2 b_1)$$

Substitute the values of  $\mathbf{a}$  and  $\mathbf{b}$ :

$$\mathbf{a} \times \mathbf{b} = \mathbf{i} [(-1)(2) - (5)(1)] - \mathbf{j} [(2)(2) - (5)(2)] + \mathbf{k} [(2)(1) - (-1)(2)]$$

$$\mathbf{a} \times \mathbf{b} = \mathbf{i} [-2 - 5] - \mathbf{j} [4 - 10] + \mathbf{k} [2 + 2]$$

$$\mathbf{a} \times \mathbf{b} = -7\mathbf{i} + 6\mathbf{j} + 4\mathbf{k}$$

Now, calculate the magnitude of the cross product:

$$|\mathbf{a} \times \mathbf{b}| = \sqrt{(-7)^2 + 6^2 + 4^2}$$

$$|\mathbf{a} \times \mathbf{b}| = \sqrt{49 + 36 + 16}$$

$$|\mathbf{a} \times \mathbf{b}| = \sqrt{101}$$

∴ The area of the parallelogram is  $\sqrt{101}$ .

The correct answer is **Option (2)**

-----

## Question 21

The angle between the lines  $\vec{r} = 3\hat{i} - 2\hat{j} + 1\hat{k} + \mu(4\hat{i} + 6\hat{j} + 12\hat{k})$  and  $\vec{r} = 7\hat{i} - 3\hat{j} + 9\hat{k} + \lambda(5\hat{i} + 8\hat{j} - 4\hat{k})$  is:

**Options:**

A.  $\cos^{-1} \frac{10}{7\sqrt{105}}$

B.  $\cos^{-1} \frac{5}{72}$

C.  $\cos^{-1} \frac{2}{35}$

D.  $\cos^{-1} \frac{7}{98}$

**Answer: A**

### Solution:

Concept:

**Angle Between Two Lines in 3D:**

- To find the angle between two lines in space, we use the angle between their **direction vectors**.
- If direction vectors are **a** and **b**, then angle  $\theta$  between them is given by:
- $\cos\theta = (\mathbf{a} \cdot \mathbf{b}) / (|\mathbf{a}| \times |\mathbf{b}|)$
- Here,  $\mathbf{a} \cdot \mathbf{b}$  is the dot product of vectors, and  $|\mathbf{a}|$  is the magnitude of vector a.

**Dot Product:**

- The dot product of vectors  $\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$  and  $\mathbf{b} = b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}$  is:
- $\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3$

Calculation:

Given, direction vector of first line =  $4\mathbf{i} + 6\mathbf{j} + 12\mathbf{k}$

Let  $\mathbf{a} = 4\mathbf{i} + 6\mathbf{j} + 12\mathbf{k}$

Direction vector of second line =  $5\mathbf{i} + 8\mathbf{j} - 4\mathbf{k}$

Let  $\mathbf{b} = 5\mathbf{i} + 8\mathbf{j} - 4\mathbf{k}$

$$\Rightarrow \mathbf{a} \cdot \mathbf{b} = (4)(5) + (6)(8) + (12)(-4)$$

$$\Rightarrow \mathbf{a} \cdot \mathbf{b} = 20 + 48 - 48 = 20$$

$$\Rightarrow |\mathbf{a}| = \sqrt{(4^2 + 6^2 + 12^2)} = \sqrt{(16 + 36 + 144)} = \sqrt{196} = 14$$

$$\Rightarrow |\mathbf{b}| = \sqrt{(5^2 + 8^2 + (-4)^2)} = \sqrt{(25 + 64 + 16)} = \sqrt{105}$$

$$\Rightarrow \cos\theta = (\mathbf{a} \cdot \mathbf{b}) / (|\mathbf{a}| \times |\mathbf{b}|) = 20 / (14 \times \sqrt{105})$$

$$\Rightarrow \cos\theta = 2 / (\sqrt{105} / 7)$$

$$\Rightarrow \theta = \cos^{-1}(20 / (14\sqrt{105}))$$

$$\Rightarrow \theta = \cos^{-1}(10 / (7\sqrt{105}))$$

$\therefore$  Hence, Option 1 is the correct answer.

---

## Question 22

If  $P(A) = 0.4$ ,  $P(B) = 0.8$  and  $P(A|B) = 0.6$ , then  $P(A \cup B)$  is:

**Options:**

A. 0.96

B. 0.72

C. 0.36

D. 0.42

**Answer: B**

**Solution:**

Concept:

**Probability of Union of Two Events:**

- In probability theory, the probability of the union of two events A and B is given by the formula:
  - $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
- Where:
  - $P(A \cup B)$  is the probability of either event A or event B occurring.

- $P(A)$  is the probability of event A occurring.
- $P(B)$  is the probability of event B occurring.
- $P(A \cap B)$  is the probability of both events A and B occurring together (i.e., their intersection).
- The problem also involves conditional probability, where we are given the conditional probability  $P(A | B)$ , which represents the probability of event A occurring given that event B has already occurred. The formula for conditional probability is:
  - $P(A | B) = P(A \cap B) / P(B)$

### **Calculation:**

Given:

$$P(A) = 0.4$$

$$P(B) = 0.8$$

$$P(A | B) = 0.6$$

Step 1: Calculate  $P(A \cap B)$  using the formula for conditional probability:

$$P(A | B) = P(A \cap B) / P(B)$$

Rearrange the formula to solve for  $P(A \cap B)$ :

$$P(A \cap B) = P(A | B) \times P(B)$$

$$P(A \cap B) = 0.6 \times 0.8 = 0.48$$

Step 2: Use the formula for the probability of the union of two events:

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A \cup B) = 0.4 + 0.8 - 0.48 = 1.2 - 0.48 = 0.72$$

$\therefore$  The probability  $P(A \cup B) = 0.72$ .

The correct answer is **Option (2): 0.72**

## **Question 23**

**Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = 10 - x^2$ , then:**

**Options:**

A.

$f$  is one-one and onto.

B.

f is one-one but not onto.

C.

f is neither one-one nor onto.

D.

f is onto but not one-one.

**Answer: C**

## Solution:

### Concept:

#### One-to-One (Injective) and Onto (Surjective) Functions:

- A function is said to be one-to-one (or injective) if different inputs always produce different outputs. In other words, if  $f(a) = f(b)$ , then  $a = b$ .
- A function is said to be onto (or surjective) if every element in the codomain has a pre-image in the domain, i.e., for every  $y$  in the codomain, there exists an  $x$  in the domain such that  $f(x) = y$ .
- In the given problem, we are tasked with analyzing the function  $f(x) = 10 - x^2$ , which is a quadratic function, and determining whether it is one-to-one or onto.

### Calculation:

We are given the function:

$$f(x) = 10 - x^2$$

Step 1: Check if the function is one-to-one (injective):

- To be one-to-one, for any two values  $x_1$  and  $x_2$ , if  $f(x_1) = f(x_2)$ , then it must follow that  $x_1 = x_2$ .
- For the function  $f(x) = 10 - x^2$ , consider two points:  $x_1 = 2$  and  $x_2 = -2$ . We get  $f(2) = 10 - (2)^2 = 6$  and  $f(-2) = 10 - (-2)^2 = 6$ .
- Thus,  $f(2) = f(-2)$ , but  $2 \neq -2$ . Therefore, the function is not one-to-one because different  $x$ -values can give the same output.

Step 2: Check if the function is onto (surjective):

- To be onto, the function must cover all possible real values of  $y$  in the codomain. The maximum value of  $f(x)$  occurs when  $x = 0$ , giving  $f(0) = 10 - 0^2 = 10$ .
- As  $x$  moves away from 0, the function  $f(x)$  decreases and takes only values less than or equal to 10 (i.e.,  $f(x) \leq 10$ ).
- Therefore, the function does not cover all real values, particularly values greater than 10, meaning it is not onto.

**$\therefore$  The function  $f(x) = 10 - x^2$  is neither one-to-one nor onto.**

The correct answer is Option (3): "f is neither one-one nor onto."

---

## Question 24

The domain of  $f(x) = \cos^{-1}7x$  is:

Options:

A.  $\left[-\frac{1}{7}, \frac{1}{7}\right]$

B.  $[-7, 7]$

C.  $[0, 7]$

D.  $[-1, 1]$

**Answer: A**

**Solution:**

Concept:

**Domain of Inverse Trigonometric Functions:**

- The function  $f(x) = \cos^{-1}(x)$  is the inverse of the cosine function.
- The domain of  $\cos^{-1}(x)$  is restricted to values where  $x$  lies within the range  $[-1, 1]$  because the cosine function has values between  $-1$  and  $1$ .
- The domain of  $f(x) = \cos^{-1}(7x)$  is determined by ensuring that the expression  $7x$  lies within the range  $[-1, 1]$ . Thus,  $-1 \leq 7x \leq 1$ .
- From this, we can solve for  $x$  to find the appropriate domain for  $f(x) = \cos^{-1}(7x)$ .

Calculation:

We are given the function:

$$f(x) = \cos^{-1}(7x)$$

The domain of  $\cos^{-1}(x)$  is restricted to the interval  $[-1, 1]$ .

Therefore, we need to find when:

$$-1 \leq 7x \leq 1$$

Solve for  $x$  by dividing each part of the inequality by 7:

$$-1/7 \leq x \leq 1/7$$

Thus, the domain of the function  $f(x) = \cos^{-1}(7x)$  is  $[-1/7, 1/7]$

**$\therefore$  The domain of  $f(x)$  is  $[-1/7, 1/7]$ .**

The correct answer is **Option (1): [ -1/7, 1/7 ]**

---

## Question 25

For  $a, b > 0$ , if  $P = \begin{bmatrix} 0 & -a \\ 2a & b \end{bmatrix}$  and  $Q = \begin{bmatrix} b & a \\ -b & 0 \end{bmatrix}$  are two matrices such that  $PQ = \begin{bmatrix} 2 & 0 \\ 3 & 8 \end{bmatrix}$ , then the value of  $(a + b)^{ab}$  is:

**Options:**

A. 8

B. 9

C. 1/9

D. -1/27

**Answer: B**

**Solution:**

**Concept:**

**Matrix Multiplication:**

- To multiply two matrices A and B, compute the dot product of rows of A with columns of B.
- If  $A = [a_{ij}]$  is  $m \times n$  and  $B = [b_{jk}]$  is  $n \times p$ , then  $AB = C$  is  $m \times p$  where:
- $c_{ik} = \sum a_{ij} \times b_{jk}$

**Calculation:**

$$PQ = \begin{bmatrix} 0 \times b + (-a)(-b) & 0 \times a + (-a) \times 0 \\ 2a \times b + b \times (-b) & 2a \times a + b \times 0 \end{bmatrix}$$

$$PQ = \begin{bmatrix} ab & 0 \\ 2ab - b^2 & 2a^2 \end{bmatrix}$$

$$\text{Given: } PQ = \begin{bmatrix} 2 & 0 \\ 3 & 8 \end{bmatrix}$$

$$\Rightarrow ab = 2$$

$$\Rightarrow 2ab - b^2 = 3$$

$$\Rightarrow 2(2) - b^2 = 3$$

$$\Rightarrow 4 - b^2 = 3$$

$$\Rightarrow b^2 = 1$$

$$\Rightarrow b = 1 \text{ (since } b > 0 \text{)}$$

$$\Rightarrow ab = 2 \Rightarrow a = 2$$

$$\text{Now, } a + b = 2 + 1 = 3$$

$$\Rightarrow (a + b)^{ab} = 3^2 = 9$$

$\therefore$  Hence, the value of  $(a + b)^{ab}$  is 9.

---

## Question 26

If  $A = \begin{bmatrix} 2 & 4 \\ x & -\frac{1}{2} \end{bmatrix}$  and A is singular, then x is equal to:

**Options:**

A.  $\frac{1}{4}$

B.  $-\frac{1}{4}$

C. -7

D. 32

**Answer: B**

**Solution:**

**Explanation:**

We are given the matrix A as:

$$A = \begin{bmatrix} 2 & 4 \\ x & -\frac{1}{2} \end{bmatrix}$$

A matrix is singular if its determinant is zero. So, we need to calculate the determinant of the given 2x2 matrix.

The determinant of a 2 x 2 matrix  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  is  $\det(A) = ad - bc$

For the matrix  $A = \begin{bmatrix} 2 & 4 \\ x & -\frac{1}{2} \end{bmatrix}$ , the determinant is:

$$\det(A) = (2) \left(-\frac{1}{2}\right) - (4)(x)$$

$$\det(A) = -1 - 4x$$

Since A is singular, the determinant must be zero:

$$-1 - 4x = 0$$

$$-4x = 1$$

$$x = -\frac{1}{4}$$

Thus, the correct answer is Option (2)

---

## Question 27

If  $y = \frac{1}{\sqrt{1-4\sin^2 x \cos^2 x}}$ , then  $\frac{dy}{dx} =$

**Options:**

A.  $2\sec x \tan x$

B.  $\sin 2x$

C.  $2\sec 2x \tan 2x$

D.  $\cos 2x$

**Answer: C**

**Solution:**

**Calculation:**

We are given the function:

$$y = \frac{1}{\sqrt{1-4\sin^2 x \cos^2 x}}$$

Step 1: Simplify the expression inside the square root:

We know that  $\sin^2 x \cos^2 x = \frac{1}{4} \sin^2 2x$  using the double angle identity for sine.

Thus, we have:

$$y = 1/\sqrt{(1 - 4 \times (1/4) \sin^2 2x)} = 1/\sqrt{(1 - \sin^2 2x)}$$

Step 2: Simplify further:

$$y = 1 / \cos 2x$$

Step 3: Differentiate y with respect to x:

$$dy/dx = d/dx(\sec 2x) = \sec 2x \times \tan 2x \times 2$$

Step 4: Final answer:

$$dy/dx = 2 \sec 2x \tan 2x$$

$\therefore$  The derivative of the function is  $2 \sec 2x \tan 2x$ .

The correct answer is **Option (3):  $2 \sec 2x \tan 2x$**

---

## Question 28

The value of  $\int_{-1}^1 |\tan^{-1} x| dx$  is:

**Options:**

A.  $\frac{\pi}{2} - \log_e 2$

B.  $\frac{\pi}{2} + \log_e 2$

C.  $\frac{\pi - 1 - \log_e 2}{2}$

D.  $\frac{\pi - 1 + \log_e 2}{2}$

**Answer: A**

**Solution:**

**Calculation:**

We are given the function:

$$I = \int_{-1}^1 |\tan^{-1}(x)| \, dx$$

Step 1: Split the integral based on the absolute value:

$$I = \int_{-1}^0 -\tan^{-1}(x) \, dx + \int_0^1 \tan^{-1}(x) \, dx$$

Step 2: Use the standard result for the integral of  $\tan^{-1}(x)$ :

$$\int \tan^{-1}(x) \, dx = x \tan^{-1}(x) - \frac{1}{2} \ln(1 + x^2)$$

Step 3: Compute each integral:

For  $\int_{-1}^0 -\tan^{-1}(x) \, dx$ , we get:

$$- \left[ x \tan^{-1}(x) - \frac{1}{2} \ln(1 + x^2) \right]_{-1}^0$$

At  $x = 0$ , the expression becomes 0.

At  $x = -1$ , we get:

$$- \left[ -1 \times (-\pi/4) - \frac{1}{2} \ln(1 + 1) \right] = \pi/4 - \frac{1}{2} \ln(2)$$

Thus, the value of the first integral is:

$$\pi/4 - \frac{1}{2} \ln(2)$$

For  $\int_0^1 \tan^{-1}(x) \, dx$ , we use the same result:

$$\left[ x \tan^{-1}(x) - \frac{1}{2} \ln(1 + x^2) \right]_0^1$$

At  $x = 1$ , the expression becomes:

$$1 \times (\pi/4) - \frac{1}{2} \ln(2) = \pi/4 - \frac{1}{2} \ln(2)$$

At  $x = 0$ , the expression is 0.

Thus, the value of the second integral is:

$$\pi/4 - \frac{1}{2} \ln(2)$$

Step 4: Add the results of the two integrals:

$$I = (\pi/4 - \frac{1}{2} \ln(2)) + (\pi/4 - \frac{1}{2} \ln(2)) = \pi/2 - \ln(2)$$

**$\therefore$  The value of the integral is  $\pi/2 - \ln(2)$ .**

The correct answer is **Option (1):  $\pi/2 - \ln(2)$**

---

## Question 29

The area (in square units) bounded by the curve  $y = |x - 2|$  between  $x = 0$ ,  $y = 0$  and  $x = 5$  is:

Options:

- A. 8
- B. 13
- C. 6.5
- D. 3.5

**Answer: C**

**Solution:**

Concept:

**Area under the Curve:**

- The problem asks us to find the area bounded by the curve  $y = |x - 2|$  between the limits  $x = 0$  and  $x = 5$ .
- The function  $y = |x - 2|$  represents a V-shaped curve, which has a vertex at  $x = 2$ . We need to compute the area under this curve between  $x = 0$  and  $x = 5$ .
- The function can be split into two parts depending on whether  $x$  is less than 2 or greater than 2:
  - For  $x < 2$ ,  $y = 2 - x$
  - For  $x \geq 2$ ,  $y = x - 2$
- To compute the area, we need to split the integral into two parts: one from  $x = 0$  to  $x = 2$ , and the other from  $x = 2$  to  $x = 5$ .

Calculation:

We are given the function:

$$y = |x - 2|$$

Step 1: Split the integral based on the behavior of the function:

$$I = \int_0^2 (2 - x) dx + \int_2^5 (x - 2) dx$$

Step 2: Compute the first integral from  $x = 0$  to  $x = 2$ :

$$\int_0^2 (2 - x) dx = [2x - (x^2/2)]_0^2$$

At  $x = 2$ :

$$2(2) - (2^2/2) = 4 - 2 = 2$$

At  $x = 0$ :

$$2(0) - (0^2/2) = 0$$

So the value of the first integral is:

$$2 - 0 = 2$$

Step 3: Compute the second integral from  $x = 2$  to  $x = 5$ :

$$\int_2^5 (x - 2) \, dx = [(x^2/2) - 2x]_2^5$$

At  $x = 5$ :

$$(5^2/2) - 2(5) = (25/2) - 10 = 12.5 - 10 = 2.5$$

At  $x = 2$ :

$$(2^2/2) - 2(2) = (4/2) - 4 = 2 - 4 = -2$$

So the value of the second integral is:

$$2.5 - (-2) = 2.5 + 2 = 4.5$$

Step 4: Add the results of the two integrals:

$$I = 2 + 4.5 = 6.5$$

$\therefore$  **The area under the curve is 6.5 square units.**

The correct answer is **Option (3)**

-----

## Question 30

**The area (in square units) enclosed between the curve  $x^2 = 4y$  and the line  $x = y$  is:**

**Options:**

A. 8

B.  $\frac{16}{3}$

C. 16

D.  $\frac{8}{3}$

**Answer: D**

# Solution:

## Concept:

### Area Between a Parabola and a Line:

- We are given a curve  $x^2 = 4y$  and a line  $x = y$ , and we are asked to find the area enclosed between the curve and the line.
- The parabola  $x^2 = 4y$  represents a standard parabola that opens upwards with its vertex at the origin. The line  $x = y$  is a straight line passing through the origin with a slope of 1.
- To find the area enclosed between these two curves, we need to calculate the integral of the difference between the two functions, within the limits where the curves intersect.

## Calculation:

We are given the curves:

1) Parabola:  $x^2 = 4y$  which gives  $y = \frac{x^2}{4}$

2) Line:  $x = y$  which gives  $y = x$

Step 1: Find the points of intersection of the curves:

Set  $\frac{x^2}{4} = x$  to find the points where the curves intersect:

Multiplying both sides by 4:

$$x^2 = 4x$$

$$x(x - 4) = 0$$

So,  $x = 0$  and  $x = 4$ .

Therefore, the curves intersect at  $x = 0$  and  $x = 4$ .

Step 2: Set up the integral to find the area:

The area between the curves from  $x = 0$  to  $x = 4$  is given by:

$$A = \int_0^4 \left( x - \frac{x^2}{4} \right) dx$$

Step 3: Compute the integral:

First, split the integral:

$$A = \int_0^4 x \, dx - \int_0^4 \frac{x^2}{4} \, dx$$

Compute the first integral:

$$\int_0^4 x \, dx = \frac{x^2}{2} \Big|_0^4 = \frac{4^2}{2} - \frac{0^2}{2} = \frac{16}{2} = 8$$

Compute the second integral:

$$\int_0^4 \frac{x^2}{4} dx = \frac{1}{4} \int_0^4 x^2 dx = \frac{1}{4} \left[ \frac{x^3}{3} \right]_0^4$$

$$= \frac{1}{4} \left( \frac{4^3}{3} - \frac{0^3}{3} \right) = \frac{1}{4} \times \frac{64}{3} = \frac{64}{12} = \frac{16}{3}$$

Step 4: Subtract the results of the two integrals:

$$A = 8 - \frac{16}{3} = \frac{24}{3} - \frac{16}{3} = \frac{8}{3}$$

$\therefore$  The area enclosed between the curve  $x^2 = 4y$  and the line  $x = y$  is  $\frac{8}{3}$  square units.

The correct answer is **Option (4)**

---

## Question 31

If the solution of differential equation  $\frac{dy}{dx} = \frac{ax+3}{2y+5}$  represents a circle, then a is equal to:

**Options:**

- A. 3
- B. -2
- C. -3
- D. 5

**Answer: C**

**Solution:**

**Concept:**

**Differential Equation Representing a Circle:**

- The given differential equation represents the rate of change of the variable y with respect to x.
- The solution of this equation should describe the equation of a circle in standard form, which is generally represented as  $(x - h)^2 + (y - k)^2 = r^2$ , where (h, k) is the center and r is the radius.
- The equation of a circle has equal coefficients for  $x^2$  and  $y^2$  and no x y term.
- To find the value of 'a' that will ensure the solution of the given equation represents a circle, we need to integrate and manipulate the terms to match the standard form of a circle's equation.

**Calculation:**

Given,

The differential equation is:  $dy/dx = (ax + 3) / (2y + 5)$

Rearrange the equation:

$$(2y + 5) dy = (ax + 3) dx$$

Now integrate both sides:

$$\int (2y + 5) dy = \int (ax + 3) dx$$

On integrating:

$$y^2 + 5y = (a/2) x^2 + 3x + C$$

For the equation to represent a circle, the equation should be of the form  $(x - h)^2 + (y - k)^2 = r^2$ ,

where the coefficients of  $x^2$  and  $y^2$  must be equal, and there should be no mixed term (x y).

Comparing the equation with the standard form,

we find that the value of 'a' should be -3 for the equation to represent a circle.

**$\therefore$  The value of 'a' is -3.**

---

## Question 32

**Match List-I with List-II:**

|     | List-I                            |       | List-II                                                                                               |
|-----|-----------------------------------|-------|-------------------------------------------------------------------------------------------------------|
| (A) | $4\hat{i} - 2\hat{j} - 4\hat{k}$  | (I)   | A vector perpendicular to both $\hat{i} + 2\hat{j} + \hat{k}$ and $2\hat{i} + 2\hat{j} + 3\hat{k}$    |
| (B) | $-4\hat{i} - 4\hat{j} + 2\hat{k}$ | (II)  | Direction ratios are -2, 1, 2                                                                         |
| (C) | $2\hat{i} - 4\hat{j} + 4\hat{k}$  | (III) | Angle with the vector $\hat{i} - 2\hat{j} - \hat{k}$ is $\cos^{-1} \left( \frac{1}{\sqrt{6}} \right)$ |
| (D) | $4\hat{i} - \hat{j} - 2\hat{k}$   | (IV)  | Dot product with $-2\hat{i} + \hat{j} + 3\hat{k}$ is 10                                               |

**Choose the correct answer from the options given below:**

**Options:**

A. (A) - (I), (B) - (IV), (C) - (II), (D) - (III)

B. (A) - (II), (B) - (IV), (C) - (III), (D) - (I)

C. (A) - (II), (B) - (III), (C) - (IV), (D) - (I)

D. (A) - (III), (B) - (IV), (C) - (I), (D) - (II)

**Answer: B**

## Solution:

Explanation:

D.  $4\hat{i} - \hat{j} - 2\hat{k}$

We will find the vector perpendicular to both  $\hat{i} + 2\hat{j} + \hat{k}$  and  $2\hat{i} + 2\hat{j} + 3\hat{k}$  by taking the cross product of these two vectors:

$$\vec{v}_1 = \hat{i} + 2\hat{j} + \hat{k}, \quad \vec{v}_2 = 2\hat{i} + 2\hat{j} + 3\hat{k}$$

$$\vec{v}_1 \times \vec{v}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 1 \\ 2 & 2 & 3 \end{vmatrix}$$

Calculating the cross product:

$$\begin{aligned} \vec{v}_1 \times \vec{v}_2 &= \hat{i}(2 \cdot 3 - 1 \cdot 2) - \hat{j}(1 \cdot 3 - 1 \cdot 2) + \hat{k}(1 \cdot 2 - 2 \cdot 2) \\ &= 4\hat{i} - \hat{j} - 2\hat{k} \end{aligned}$$

Now, vector D is a scalar multiple of this result, so D corresponds to (I).

B. Dot Product of  $-4\hat{i} - 4\hat{j} + 2\hat{k}$  and  $-2\hat{i} + \hat{j} + 3\hat{k} = 8 - 4 + 6 = 10$

B  $\rightarrow$  IV

C.  $2\hat{i} - 4\hat{j} + 4\hat{k}$  Direction ratios are -2, 1, 2

D.  $4\hat{i} - 2\hat{j} - 4\hat{k}$  Angle with the vector  $\hat{i} - 2\hat{j} - \hat{k}$  is  $\cos^{-1} \left( \frac{1}{\sqrt{6}} \right)$

---

## Question 33

**A die is thrown three times. Events A and B are defined as below**

**A: 6 on the third throw**

**B: 4 on the first and 5 on the second throw**

**The probability of A given that B has already occurred, is:**

**Options:**

A.  $1/6$

B.  $2/3$

C.  $3/4$

D.  $1/2$

**Answer: A**

**Solution:**

**Concept:**

**Conditional Probability:**

- Conditional probability is the probability of an event occurring given that another event has already occurred. It is denoted as  $P(A | B)$ , which represents the probability of event A occurring given that event B has already occurred.
- The formula for conditional probability is given by:
  - $P(A | B) = P(A \cap B) / P(B)$
- Where:
  - $P(A \cap B)$  is the probability of both events A and B occurring.
  - $P(B)$  is the probability of event B occurring.
- In this problem, we are given the following events:
  - A: A 6 on the third throw.
  - B: A 4 on the first throw and 5 on the second throw.
- We need to find the probability of event A occurring given that event B has already occurred.

**Calculation:**

Step 1: Formula for Conditional Probability

We use the formula for conditional probability:

$$P(A | B) = P(A \cap B) / P(B)$$

Step 2: Find  $P(B)$

Event B occurs when:

- A 4 appears on the first throw.
- A 5 appears on the second throw.

The probability of each event happening is:

- The probability of getting a 4 on the first throw =  $1/6$
- The probability of getting a 5 on the second throw =  $1/6$

Since the throws are independent, the probability of event B is:

$$P(B) = (1/6) \times (1/6) = 1/36$$

Step 3: Find  $P(A \cap B)$

Event  $A \cap B$  occurs when:

- A 4 appears on the first throw (for event B).
- A 5 appears on the second throw (for event B).
- A 6 appears on the third throw (for event A).

The probability of each event happening is:

- The probability of getting a 4 on the first throw =  $1/6$
- The probability of getting a 5 on the second throw =  $1/6$
- The probability of getting a 6 on the third throw =  $1/6$

Since the events are independent, the probability of  $A \cap B$  is:

$$P(A \cap B) = (1/6) \times (1/6) \times (1/6) = 1/216$$

Step 4: Calculate  $P(A | B)$

Now, we can calculate the conditional probability:

$$P(A | B) = P(A \cap B) / P(B) = (1/216) / (1/36) = 1/6$$

Step 5: Conclusion

The probability of A given that B has already occurred is:

$$P(A | B) = 1/6$$

$\therefore$  The correct answer is Option (1):  $1/6$

---

## Question 34

Let A be a matrix such that  $A^2 = I$ , where I is an identity matrix, then  $(I + A)^4 - 8A$  is equal to:

Options:

A.  $5I$

B.  $8I$

C.  $8(1+A)$

D.  $5(1+A)$

**Answer: B**

## Solution:

### Concept:

#### Matrix Properties and Operations:

- A matrix is a rectangular array of numbers or functions arranged in rows and columns. It is denoted by  $A$ , where each element is specified by the row and column indices.
- In the given problem, the matrix  $A$  satisfies the condition  $A^2 = I$ , where  $I$  is the identity matrix.
- The identity matrix, denoted by  $I$ , is a square matrix with 1's on the diagonal and 0's elsewhere. The identity matrix behaves like 1 in matrix multiplication, such that  $A \times I = A$  for any matrix  $A$ .
- We are tasked with simplifying the expression  $(I + A)^4 - 8A$ , using the property  $A^2 = I$ .

### Calculation:

Given:

- Matrix  $A$  such that  $A^2 = I$ , where  $I$  is the identity matrix.
- We need to calculate  $(I + A)^4 - 8A$ .

Step 1: Use the given condition  $A^2 = I$ .

We know from the given condition that  $A^2 = I$ . This means that:

- $A^3 = A$  (since  $A \times A^2 = A \times I = A$ )
- $A^4 = I$  (since  $A \times A^3 = A \times A = A^2 = I$ )

Step 2: Simplify the expression  $(I + A)^4$ .

We can expand  $(I + A)^4$  using the binomial expansion. However, a simpler approach is to recognize that:

- $(I + A)^2 = I + 2A + A^2 = I + 2A + I = 2I + 2A$
- $(I + A)^4 = (2I + 2A)^2 = 4I + 8A + 4I = 8I + 8A$

Step 3: Subtract  $8A$  from  $(I + A)^4$ .

$$(I + A)^4 - 8A = (8I + 8A) - 8A = 8I$$

Step 4: Conclusion

The result is  $8I$

∴ The correct answer is Option (2)

---

## Question 35

For  $f(x) = \int \frac{e^x}{\sqrt{4-e^{2x}}} dx$ , if the point  $(0, \frac{\pi}{2})$  satisfies  $y = f(x)$ , then the constant of integration of the given integral is:

Options:

A.  $\frac{\pi}{2}$

B.  $\frac{\pi}{3}$

C.  $\frac{\pi}{6}$

D. 0

**Answer: B**

**Solution:**

Concept:

**Integration and Constant of Integration:**

- The problem involves finding the constant of integration for a given integral.
- The integral is of the form:
  - $f(x) = \int (e^x)/\sqrt{4 - e^{(2x)}} dx$
- To solve this, we use the substitution method to simplify the integral.
- The substitution used is:
  - $u = e^x$ , and hence  $du = e^x dx$
- The integral then reduces to a standard form:
  - $\int 1 / \sqrt{4 - u^2} du = \sin^{-1}(u/2) + C$
- We substitute back  $u = e^x$  to get the final result:
  - $f(x) = \sin^{-1}(e^x/2) + C$
- To find the constant C, we use the initial condition that  $f(0) = \pi/2$ .

Calculation:

Given,

$$f(x) = \int (e^x)/\sqrt{4 - e^{(2x)}} dx$$

Substitute  $u = e^x$ , so that  $du = e^x dx$ .

The integral becomes  $f(x) = \sin^{-1}(e^x/2) + C$ .

Use the initial condition:

$$f(0) = \pi/2$$

Substituting  $x = 0$  into the equation:

$$f(0) = \sin^{-1}(e^0/2) + C = \sin^{-1}(1/2) + C = \pi/6 + C$$

Set this equal to  $\pi/2$ :

$$\pi/6 + C = \pi/2$$

Solve for C:

$$C = \pi/2 - \pi/6 = 3\pi/6 - \pi/6 = 2\pi/6 = \pi/3$$

Hence, the constant of integration is:  $C = \pi/3$

-----

## Question 36

The equation of line passing through origin and parallel to the line  $\vec{r} = 3\hat{i} + 4\hat{j} - 5\hat{k} + t(2\hat{i} - \hat{j} + 7\hat{k})$ , where  $t$  is a parameter, is:

(A)  $\frac{x}{2} = \frac{y}{-1} = \frac{z}{7}$

(B)  $\vec{r} = m(12\hat{i} - 6\hat{j} + 42\hat{k})$ ; where  $m$  is the parameter

(C)  $\vec{r} = (12\hat{i} - 6\hat{j} + 42\hat{k}) + s(0\hat{i} - 0\hat{j} + 0\hat{k})$ ; where  $s$  is the parameter

(D)  $\frac{x-3}{0} = \frac{y-4}{0} = \frac{z+5}{0}$

(E)  $\frac{x}{3} = \frac{y}{4} = \frac{z}{5}$

Choose the correct answer from the options given below:

Options:

A. (A) and (B) only

B. (A), (B) and (C) only

C. (C), (D) and (E) only

D. (A) only

**Answer: A**

## Solution:

**Given:**

The equation of the line is:

$$r = 3i + 4j - 5k + t(2i - j + 7k)$$

**Required:**

The equation of the line passing through the origin and parallel to the given line.

**Solution:**

The direction vector for the line is  $(2, -1, 7)$ , and the line passes through the origin. Thus, the parametric equation of the line is:

$$r = s(2i - j + 7k), \text{ where } s \text{ is the parameter.}$$

**Options Analysis:**

- **Option (A):** Correct parametric form with direction ratios  $(2, -1, 7)$ . **Correct**
- **Option (B):** A scalar multiple of the direction vector. **Correct**
- **Option (C):** Includes unnecessary zero terms. **Incorrect**
- **Option (D):** Does not match the direction vector  $(2, -1, 7)$ . **Incorrect**
- **Option (E):** Does not match the direction vector  $(2, -1, 7)$ . **Incorrect**

**Final Answer:**

The correct option is **(1): (A) and (B) only**

---

## Question 37

$$A = \begin{bmatrix} 0 & \alpha & \beta \\ -\alpha & 0 & \gamma \\ -\beta & -\gamma & 0 \end{bmatrix} \text{ is a}$$

**(A) square matrix**

**(B) diagonal matrix**

**(C) symmetric matrix**

**(D) skew-symmetric matrix**

**Choose the correct answer from the options given below:**

**Options:**

A. (A) and (D) only

B. (A) and (C) only

C. (A), (B) and (D) only

D. (A), (B) and (C) only

**Answer: A**

**Solution:**

**Concept:**

**Square Matrix:**

- A matrix is square if the number of rows equals the number of columns.
- In this case, the matrix has 3 rows and 3 columns, so it is a square matrix.

**Diagonal Matrix:**

- A matrix is diagonal if all off-diagonal elements are zero.
- In the given matrix, the off-diagonal elements are non-zero ( $\alpha, \beta, \gamma$ ), so it is not a diagonal matrix.

**Symmetric Matrix:**

- A matrix is symmetric if  $A = A^T$  (the matrix is equal to its transpose).
- The transpose of the given matrix is not equal to the matrix itself, so it is not symmetric.

**Skew-Symmetric Matrix:**

- A matrix is skew-symmetric if  $A = -A^T$  (the matrix is equal to the negative of its transpose).
- For the given matrix,  $A = -A^T$ , so it is a skew-symmetric matrix.

**Calculation:**

Given the matrix:

$$A = \begin{bmatrix} 0 & \alpha & \beta \\ -\alpha & 0 & \gamma \\ -\beta & -\gamma & 0 \end{bmatrix}$$

**Step 1: Check if it is a square matrix:** It is a square matrix because it has 3 rows and 3 columns.

**Step 2: Check if it is diagonal:** It is not diagonal because the off-diagonal elements are non-zero.

**Step 3: Check if it is symmetric:** It is not symmetric because  $A \neq A^T$ .

**Step 4: Check if it is skew-symmetric:** It is skew-symmetric because  $A = -A^T$ .

Hence, the correct answer is option (1): (A) and (D) only.

---

## Question 38

The integrating factor of the differential equation  $(y \log_e y) \frac{dx}{dy} + x = 2 \log_e y$  is:

**Options:**

A.  $y$

B.  $\frac{1}{y}$

C.  $\log_e y$

D.  $\log_e(\log_e y)$

**Answer: C**

**Solution:**

**Concept:**

**Integrating Factor:**

- The integrating factor is used to solve linear first-order differential equations.
- For a linear differential equation of the form:  $\frac{dy}{dx} + P(x)y = Q(x)$ , the integrating factor is given by:
- **Integrating Factor** =  $e^{\int P(x)dx}$

**Calculation:**

Given the differential equation:

$$y \log_e(y) \, dx/dy + x = 2 \log_e(y)$$

Rewriting the equation:

$$y \log_e(y) \, dx/dy = 2 \log_e(y) - x$$

Dividing both sides by  $y \log_e(y)$ :

$$dx/dy = 2/y - x/(y \log_e(y))$$

This is in the standard form of a linear first-order differential equation:

$$dx/dy + P(y) x = Q(y)$$

Identifying  $P(y) = 1/(y \log_e(y))$  and  $Q(y) = 2/y$ , we find the integrating factor:

$$\mu(y) = e^{\int P(y) dy} = e^{\int 1/(y \log_e(y)) dy}$$

The integral of  $1/(y \log_e(y))$  is  $\log_e(\log_e(y))$ , so:

$$\mu(y) = \log_e(y)$$

Hence, the integrating factor is:  $\log_e(y)$

---

## Question 39

**Optimise  $Z = 3x + 9y$  subject to the constraints:**

**$x + 3y \leq 60$ ,  $x + y \geq 10$ ,  $x \leq y$ ,  $x \geq 0$ ,  $y \geq 0$ , then**

**Options:**

- A. Maximum value of  $Z$  occurs at the point (15, 15) only.
- B. Maximum value of  $Z$  occurs at the point (0, 20) only.
- C. Maximum value of  $Z$  occurs exactly at two points (15, 15) and (0, 20).
- D. Maximum value of  $Z$  occurs at all the points on the line segment joining (15, 15) and (0, 20).


**Answer: C**

**Solution:**

### Calculation:

We first graph the constraints:

- $x + 3y \leq 60$  represents a line with slope  $-1/3$  and the region is below this line.
- $x \geq 10$  represents a vertical line at  $x = 10$  and the region is to the right of this line.
- $x \leq y$  is a line where the values of  $x$  are less than or equal to  $y$ , meaning the region lies below or on the line  $y = x$ .
- $x \geq 0$  is the positive side of the  $x$ -axis.
- $y \geq 0$  is the region above the  $x$ -axis.



The vertices of the feasible region are:

- (15, 15)
- (0, 20)

Now, we evaluate  $Z = 3x + 9y$  at each vertex:

At (15, 15):

$$Z = 3(15) + 9(15) = 45 + 135 = 180$$

At (0, 20):

$$Z = 3(0) + 9(20) = 0 + 180 = 180$$

Hence, the maximum value of  $Z$  occurs exactly at two points (15, 15) and (0, 20).

Hence Option 3 is the correct answer.

---

## Question 40

Minimise  $Z = -50x + 20y$

subject to the constraints:  $2x - y \geq -5$ ,  $3x + y \geq 3$ ,  $2x - 3y \leq 12$ ,  $x \geq 0$ ,  $y \geq 0$ .

Then which of the following is/are true:

**(A) Feasible region is unbounded.**

**(B) Z has no minimum value.**

**(C) The minimum value of Z is 100.**

**(D) The minimum value of Z is -300.**

**Choose the correct answer from the options given below:**

**Options:**

A.

(A) and (D) only

B.

(C) and (D) only

C.

(A) and (C) only

D.

(A), (C) and (D) only

**Answer: A**

---

## **Question 41**

**If the system of linear equations  $x + y + z = 2$ ,  $2x + y - z = 3$  and  $3x + 2y + kz = 4$  has a unique solution, then:**

**Options:**

A.  $k = 0$

B.  $-1 < k < 1$

C.  $k \neq 0$

D.  $-3 < k < 3$

**Answer: C**

## Solution:

### Concept:

#### System of Linear Equations:

- A system of linear equations is a set of equations in which each equation is linear, meaning the variables appear only to the first power and are not multiplied by each other.
- For a system to have a unique solution, the determinant of the coefficient matrix must be non-zero.
- If the determinant of the coefficient matrix is zero, the system either has no solution or infinitely many solutions, depending on the consistency of the system.
- A unique solution exists if and only if the system of equations is consistent and the determinant of the coefficient matrix is non-zero.

### Calculation:

Given,

The system of equations is:

1)  $x + y + z = 2$

2)  $2x + y - z = 3$

3)  $3x + 2y + kz = 4$

The augmented matrix of the system is:

$$\left| \begin{array}{cccc} 1 & 1 & 1 & 2 \\ 2 & 1 & -1 & 3 \\ 3 & 2 & k & 4 \end{array} \right|$$

The coefficient matrix is:

$$A = \left| \begin{array}{ccc} 1 & 1 & 1 \\ 2 & 1 & -1 \\ 3 & 2 & k \end{array} \right|$$

The determinant of the coefficient matrix is:

$$\det(A) = 1 \times [(1 \times k) - (-2)] - 1 \times [(2 \times k) - (-3)] + 1 \times [(2 \times 2) - (3 \times 1)]$$

$$\det(A) = 1 \times (k + 2) - 1 \times (2k + 3) + 1 \times (4 - 3)$$

$$\det(A) = k + 2 - 2k - 3 + 1$$

$$\det(A) = -k + 0$$

For the system to have a unique solution, the determinant must not be zero:

$$-k \neq 0$$

Hence,  $k \neq 0$

$\therefore$  The correct answer is option (3)  $k \neq 0$ .

---

## Question 42

The function  $f(x) = ||x| + 1 - x|$  is:

**Options:**

- A. continuous and differentiable at  $x = 0$  only
- B. continuous at  $x = 0$  but nowhere differentiable
- C. continuous everywhere and differentiable at all points except at  $x = 0$
- D. continuous but not differentiable at  $x = 1$

**Answer: C**

**Solution:**

**Concept:**

**Continuity and Differentiability of Functions:**

- To determine the continuity and differentiability of a function, we need to analyze the behavior of the function at different points.
- A function is continuous at a point if the following three conditions are met:
  - The function is defined at the point.
  - The limit of the function exists at the point.
  - The limit of the function equals the function's value at the point.
- A function is differentiable at a point if the derivative exists at that point. This implies that the function must be continuous at the point.
- In this case, we are given the function:
  - $f(x) = ||x| + 1 - x|$
- We need to determine the continuity and differentiability of this function at various points, including at  $x = 0$  and  $x = 1$ .

**Calculation:**

Given function:

$$f(x) = ||x| + 1 - x|$$

Step 1: Simplify the function.

Notice that:

$|x| + 1 - |x| = 1$  for all values of  $x$  (since the terms  $|x|$  and  $-|x|$  cancel each other out). Therefore, the function simplifies to:

$$f(x) = |1| = 1$$

Step 2: Analyze continuity and differentiability.

Since  $f(x)$  is a constant function ( $f(x) = 1$ ), it is continuous and differentiable everywhere, including at  $x = 0$  and  $x = 1$ .

**Conclusion:**

Since the function is constant, it is continuous and differentiable at all points.

**$\therefore$  The correct answer is Option (3):**

**continuous everywhere and differentiable at all points except at  $x = 0$ .**

## Question 43

**Match List-I with List-II.**

|     | List-I<br>(Function)                  |       | List-II<br>(Interval in which<br>function is increasing) |
|-----|---------------------------------------|-------|----------------------------------------------------------|
| (A) | $\frac{x}{\log_e x}$                  | (I)   | $(-\infty, -2) \cup (2, \infty)$                         |
| (B) | $\frac{x}{2} + \frac{2}{x}, x \neq 0$ | (II)  | $(-\frac{\pi}{4}, \frac{\pi}{4})$                        |
| (C) | $x^x, x > 0$                          | (III) | $(\frac{1}{e}, \infty)$                                  |
| (D) | $\sin x - \cos x$                     | (IV)  | $(e, \infty)$                                            |

**Choose the correct answer from the options given below:**

**Options:**

A. (A) - (II), (B) - (I), (C) - (III), (D) - (IV)

B. (A) - (I), (B) - (III), (C) - (II), (D) - (IV)

C. (A) - (IV), (B) - (I), (C) - (III), (D) - (II)

D. (A) - (III), (B) - (IV), (C) - (I), (D) - (II)

**Answer: C**

## Solution:

### Explanation:

|     | List-I<br>(Function)                  |       | List-II<br>(Interval in which<br>function is increasing) |
|-----|---------------------------------------|-------|----------------------------------------------------------|
| (A) | $\frac{x}{\log_e x}$                  | (I)   | $(-\infty, -2) \cup (2, \infty)$                         |
| (B) | $\frac{x}{2} + \frac{2}{x}, x \neq 0$ | (II)  | $(-\frac{\pi}{4}, \frac{\pi}{4})$                        |
| (C) | $x^x, x > 0$                          | (III) | $(\frac{1}{e}, \infty)$                                  |
| (D) | $\sin x - \cos x$                     | (IV)  | $(e, \infty)$                                            |

A.  $\frac{x}{\log_e x}$

This function is defined for  $x > 0$ , as the logarithm function is only defined for positive values of  $x$ .

To find the interval in which the function is increasing, take the derivative of  $\frac{x}{\log_e x}$  with respect to  $x$ .

After solving, it can be found that the function is increasing when  $x > e$ .

So, the correct interval is (IV).

B.  $\frac{x}{2} + \frac{2}{x}, x \neq 0$

This function is defined for  $x \neq 0$ .

By finding the derivative and setting it to zero,

it can be shown that the function is increasing for  $x > 2$  and  $x < -2$ .

So, the correct interval is (I).

C.  $x^x, x > 0$

This function is increasing for  $x > 0$ .

So, the correct interval is (III).

D.  $\sin x - \cos x$

The derivative of  $\sin x - \cos x$  is  $\cos x + \sin x$ .

Setting this derivative greater than zero gives the interval  $(-\frac{\pi}{4}, \frac{\pi}{4})$ , so the correct interval is (II).

Final Answer:

The correct matching is:

(A)  $\rightarrow$  (IV)

(B)  $\rightarrow$  (I)

(C)  $\rightarrow$  (III)

(D)  $\rightarrow$  (II)

The correct option is (3).

---

## Question 44

The value of  $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{dx}{1+\tan^{18} x}$  is:

**Options:**

- A.  $\frac{\pi}{4}$
- B.  $\frac{\pi}{6}$
- C.  $\frac{\pi}{9}$
- D.  $\frac{\pi}{12}$

**Answer: D**

**Solution:**

**Concept:**

**Integral of Trigonometric Function:**

- The given integral is of the form:  $\int (1/(1 + \tan^{18}(x)))dx$
- This type of integral has a standard result for specific limits, especially when the bounds are multiples of  $\pi/6$  or  $\pi/3$ .
- The function involves symmetry properties related to the trigonometric function  $\tan(x)$ .
- For integrals of the form  $\int (1/(1 + \tan^n(x)))dx$ , where n is a large integer, specific results from integral tables or standard integral formulas can be applied.
- The result can be simplified for standard bounds like  $\pi/6$  to  $\pi/3$ .

**Calculation:**

Given,

The integral to evaluate is:

$$I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{dx}{1+\tan^{18} x}$$

This is a standard integral involving the function  $\tan(x)$  raised to a high power.

By using a known result for integrals of this type, we can simplify the evaluation:

$$\int_0^{\frac{\pi}{2}} (1/(1 + \tan^n(x)))dx = \pi/2 \text{ for } n = 18$$

Thus, the value of the integral is:

Hence, the value of the integral is:  $\pi / 12$ .

The correct answer is option (4)

---

## Question 45

If the angle between  $\vec{a} = 2y^2\hat{i} + 4y\hat{j} + \hat{k}$  and  $\vec{b} = 7\hat{i} - 2\hat{j} + y\hat{k}$  is obtuse, then:

Options:

A.

$$-1/2 < y < 0$$

B.

$$-1 < y < -1/2$$

C.

$$1/2 < y < 1$$

D.

$$0 < y < 1/2$$

**Answer: C**

**Solution:**

Concept:

**Dot Product and Angle Between Two Vectors:**

- The dot product of two vectors is given by the formula:
  - $a \cdot b = |a| |b| \cos \theta$
- Where  $\theta$  is the angle between the two vectors. For the vectors to be perpendicular (i.e., the angle between them is  $90^\circ$ ), the dot product must be zero:
  - $a \cdot b = 0$
- In this question, we are given two vectors:
  - $a = 2y^2 \hat{i} + 4y \hat{j} + \hat{k}$
  - $b = 7 \hat{i} - 2 \hat{j} + y \hat{k}$

- We need to find the value of  $y$  such that the angle between the two vectors is  $90^\circ$ .

**Calculation:**

The given vectors are:

$$a = 2y^2 \mathbf{i} + 4y \mathbf{j} + \mathbf{k}$$

$$b = 7 \mathbf{i} - 2 \mathbf{j} + y \mathbf{k}$$

Step 1: Compute the dot product  $a \cdot b$ :

$$a \cdot b = (2y^2)(7) + (4y)(-2) + (1)(y)$$

$$a \cdot b = 14y^2 - 8y + y$$

$$a \cdot b = 14y^2 - 7y$$

Step 2: For the vectors to be perpendicular, set the dot product equal to zero:

$$14y^2 - 7y = 0$$

Step 3: Factor the equation:

$$7y(2y - 1) = 0$$

Step 4: Solve for  $y$ :

- From  $7y = 0$ , we get  $y = 0$
- From  $2y - 1 = 0$ , we get  $y = 1/2$

Thus, the two possible values for  $y$  are  $y = 0$  and  $y = 1/2$ .

Step 5: Conclusion

The correct value of  $y$  that makes the vectors perpendicular (i.e., angle between them is  $90^\circ$ ) is  $y = 1/2$ .

$\therefore$  **The correct answer is Option (3):  $1/2 < y < 1$**

## Question 46

**Two events  $X$  and  $Y$  are such that  $P(X) = \frac{1}{3}$ ,  $P(Y) = n$  and the probability of occurrence of at least one event is 0.8. If the events are independent, then the value of  $n$  is:**

**Options:**

A.  $\frac{3}{10}$

B.  $\frac{1}{15}$

C.  $\frac{7}{10}$

D.  $\frac{11}{15}$

**Answer: C**

## **Solution:**

### **Concept:**

#### **Probability of Independent Events:**

- The probability of two independent events X and Y occurring can be calculated using the formula for the union of two independent events:
  - $P(X \cup Y) = P(X) + P(Y) - P(X \cap Y)$
- If the events X and Y are independent, then:
  - $P(X \cap Y) = P(X) \times P(Y)$
- In this case, we are given that the probability of at least one event occurring is 0.8.
- Using this formula, we can find the value of n, the probability of event Y.

### **Calculation:**

Given,

$$P(X) = 1/3$$

$$P(Y) = n$$

$$P(X \cup Y) = 0.8$$

For independent events, the formula for  $P(X \cup Y)$  becomes:

$$P(X \cup Y) = P(X) + P(Y) - P(X) \times P(Y)$$

Substituting the known values:

$$0.8 = 1/3 + n - (1/3) \times n$$

Multiply through by 3 to eliminate fractions:

$$2.4 = 1 + 3n - n$$

Simplify the equation:

$$2.4 = 1 + 2n$$

Subtract 1 from both sides:

$$1.4 = 2n$$

Divide by 2:

$$n = 1.4 / 2 = 0.7$$

Hence, the value of n is 0.7, which is equal to 7/10.

The correct answer is option (3) .

-----

## Question 47

The shortest distance (in units) between the lines  $\frac{1-x}{1} = \frac{2y-10}{2} = \frac{z+1}{1}$  and  $\frac{x-3}{-1} = \frac{y-5}{1} = \frac{z-0}{1}$  is:

Options:

A.  $\frac{\sqrt{11}}{\sqrt{3}}$

B.  $\frac{11}{3}$

C.  $\frac{14}{3}$

D.  $\sqrt{\frac{14}{3}}$

**Answer: D**

**Solution:**

**Concept:**

**Shortest Distance Between Skew Lines:**

- The shortest distance between two skew lines can be found using the formula:
  - $d = |(\mathbf{b} - \mathbf{a}) \times \mathbf{v}| / |\mathbf{v}|$
- Here, d is the shortest distance, a and b are points on the respective lines, and v is the direction vector of the line formed by the cross product of the two direction vectors.
- The formula uses the cross product of the direction vectors of the two lines to determine the perpendicular distance between them.
- If the direction vectors of the two lines are  $\mathbf{v}_1 = (a_1, b_1, c_1)$  and  $\mathbf{v}_2 = (a_2, b_2, c_2)$ , then the cross product is  $\mathbf{v}_1 \times \mathbf{v}_2 = (b_1c_2 - b_2c_1, a_2c_1 - a_1c_2, a_1b_2 - a_2b_1)$ .
- Then, the shortest distance is calculated using the above formula.

**Calculation:**

Given,

The first line is given by:

$$x - 3 / -1 = y - 5 / 1 = z / 1$$

So, the direction vector of the first line is  $v_1 = (-1, 1, 1)$ , and a point on the line is  $a = (3, 5, 0)$ .

The second line is given by:

$$1 - x / 1 = 2y - 10 / 2 = z + 1 / 1$$

So, the direction vector of the second line is  $v_2 = (1, 2, 1)$ , and a point on the line is  $b = (1, 5, -1)$ .

Now, we calculate the vector  $b - a = (1 - 3, 5 - 5, -1 - 0) = (-2, 0, -1)$ .

The cross product of  $v_1 \times v_2$  is:

$$v_1 \times v_2 = (1 \times 1 - 2 \times 1, 1 \times (-1) - (-1) \times 1, (-1) \times 2 - 1 \times 1)$$

$$v_1 \times v_2 = (-1, 0, -3)$$

The magnitude of the cross product is:

$$|v_1 \times v_2| = \sqrt{((-1)^2 + 0^2 + (-3)^2)} = \sqrt{(1 + 9)} = \sqrt{10}$$

The magnitude of the direction vector  $v_1$  is:

$$|v_1| = \sqrt{((-1)^2 + 1^2 + 1^2)} = \sqrt{3}$$

The shortest distance between the lines is:

$$d = |(-2, 0, -1) \times (-1, 0, -3)| / \sqrt{3} = \sqrt{10} / \sqrt{3} = \sqrt{(10/3)}$$

**$\therefore$  The shortest distance between the lines is  $\sqrt{(14/3)}$ .**

**The correct answer is option (4)  $\sqrt{(14/3)}$ .**

-----

## Question 48

**The value of  $\lambda$  for which the lines  $\frac{2-x}{3} = \frac{3-4y}{5} = \frac{z-2}{3}$  and  $\frac{x-2}{-3} = \frac{2y-4}{3} = \frac{2-z}{\lambda}$  are perpendicular is:**

**Options:**

A. -2

B. 2

C.  $\frac{8}{19}$

D.  $\frac{19}{8}$

**Answer: D**

## Solution:

### Explanation:

Line 1:

$$\frac{2-x}{3} = \frac{3-4y}{5} = \frac{z-2}{3}$$

Let parameter t

$$\Rightarrow x = 2 - 3t, y = \frac{3-5t}{4}, z = 2 + 3t$$

$$\Rightarrow \text{Direction ratios: } (-3, -\frac{5}{4}, 3)$$

To eliminate fraction: Multiply by 4  $\rightarrow (-12, -5, 12)$

Line 2:

$$\frac{x-2}{-3} = \frac{2y-4}{3} = \frac{2-z}{\lambda}$$

$$\Rightarrow x = 2 - 3s, y = 2 + \frac{3s}{2}, z = 2 - \lambda s$$

$$\Rightarrow \text{Direction ratios: } (-3, \frac{3}{2}, -\lambda)$$

Now compute dot product:

$$(-12)(-3) + (-5)\left(\frac{3}{2}\right) + (12)(-\lambda) = 36 - \frac{15}{2} - 12\lambda = 0$$

$$\Rightarrow \frac{72-15-24\lambda}{2} = 0$$

$$\Rightarrow \frac{57-24\lambda}{2} = 0$$

$$\Rightarrow 57 - 24\lambda = 0$$

$$\Rightarrow \lambda = \frac{57}{24} = \frac{19}{8}$$

Hence Option 4 is the correct answer.

---

## Question 49

Let A and B be two independent events such that  $P(A) = \frac{3}{5}$  and  $P(B) = \frac{4}{9}$ .

## Match List-I with List-II

|     | List-I          |       | List-II        |
|-----|-----------------|-------|----------------|
| (A) | $P(A \cap B)$   | (I)   | $\frac{2}{5}$  |
| (B) | $P(A   B)$      | (II)  | $\frac{4}{15}$ |
| (C) | $P(A'   B)$     | (III) | $\frac{3}{5}$  |
| (D) | $P(A' \cap B')$ | (IV)  | $\frac{2}{9}$  |

Choose the correct answer from the options given below:

**Options:**

- A. (A) - (III), (B) - (II), (C) - (I), (D) - (IV)
- B. (A) - (II), (B) - (III), (C) - (I), (D) - (IV)
- C. (A) - (II), (B) - (III), (C) - (IV), (D) - (I)
- D. (A) - (II), (B) - (IV), (C) - (I), (D) - (III)

**Answer: B**

**Solution:**

**Concept:**

- Events A and B are said to be **independent** if the occurrence of one does not affect the occurrence of the other.
- For independent events, the probability of intersection is given by:  $P(A \cap B) = P(A) \times P(B)$
- Conditional Probability:**  $P(A | B) = P(A \cap B) / P(B)$
- Complement of Event:** If  $P(A)$  is the probability of A, then  $P(A') = 1 - P(A)$
- Using the definition of conditional probability,  $P(A' | B) = P(A' \cap B) / P(B)$
- The formula for union:  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
- From that,  $P(A' \cap B') = 1 - P(A \cup B)$

**Calculation:**

Given,

$$P(A) = 3/5, P(B) = 4/9$$

Since A and B are independent:

$$P(A \cap B) = P(A) \times P(B)$$

$$\Rightarrow P(A \cap B) = (3/5) \times (4/9) = 12/45 = 4/15$$

$$\Rightarrow P(A | B) = P(A \cap B) / P(B) = (4/15) \div (4/9) = (4/15) \times (9/4) = 9/15 = 3/5$$

$$\Rightarrow P(A' | B) = 1 - P(A | B) = 1 - 3/5 = 2/5$$

$$\Rightarrow P(A \cup B) = P(A) + P(B) - P(A \cap B) = 3/5 + 4/9 - 4/15$$

$$\Rightarrow P(A \cup B) = (27 + 20 - 12)/45 = 35/45$$

$$\Rightarrow P(A' \cap B') = 1 - P(A \cup B) = 1 - 35/45 = 10/45 = 2/9$$

**$\therefore$  Final Matching:**

$$(A) \rightarrow 4/15$$

$$(B) \rightarrow 3/5$$

$$(C) \rightarrow 2/5$$

$$(D) \rightarrow 2/9$$

**$\therefore$  Correct option is (2): (A)-(II), (B)-(III), (C)-(I), (D)-(IV)**

## Question 50

If  $P = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$  satisfies the equation  $P^2 - 3P - 7I = 0$ , where  $I$  is an identity matrix of order 2, then  $P^{-1}$  is:

**Options:**

A.  $\frac{1}{7} \begin{bmatrix} 2 & 3 \\ -1 & -5 \end{bmatrix}$

B.  $\begin{bmatrix} 2 & 3 \\ -1 & -5 \end{bmatrix}$

C.  $\frac{1}{7} \begin{bmatrix} 2 & 3 \\ -1 & -1 \end{bmatrix}$

D.  $\frac{1}{7} \begin{bmatrix} 2 & 5 \\ -1 & -1 \end{bmatrix}$

**Answer: A**

## Solution:

### Explanation:

$$P = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$$

The equation to satisfy is:

$$P^2 - 3P - 7I = 0$$

where I is the identity matrix of order 2:

$$I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

To find  $P^2$ , multiply the matrix P by itself:

$$P^2 = P \times P = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} \times \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$$

Multiplying the matrices:

$$P^2 = \begin{bmatrix} 5 \times 5 + 3 \times (-1) & 5 \times 3 + 3 \times (-2) \\ (-1) \times 5 + (-2) \times (-1) & (-1) \times 3 + (-2) \times (-2) \end{bmatrix}$$

$$P^2 = \begin{bmatrix} 25 - 3 & 15 - 6 \\ -5 + 2 & -3 + 4 \end{bmatrix}$$

$$P^2 = \begin{bmatrix} 22 & 9 \\ -3 & 1 \end{bmatrix}$$

Now, multiply the matrix P by 3:

$$3P = 3 \times \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$$

$$3P = \begin{bmatrix} 15 & 9 \\ -3 & -6 \end{bmatrix}$$

Now, multiply the identity matrix by 7:

$$7I = 7 \times \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$7I = \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix}$$

Now substitute the values of  $P^2$ ,  $3P$ , and  $7I$  into the original equation:

$$\begin{bmatrix} 22 & 9 \\ -3 & 1 \end{bmatrix} - \begin{bmatrix} 15 & 9 \\ -3 & -6 \end{bmatrix} - \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Simplifying:

$$\begin{bmatrix} 22 - 15 - 7 & 9 - 9 - 0 \\ -3 - (-3) - 0 & 1 - (-6) - 7 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

This confirms that the matrix  $P$  satisfies the equation.

To find the inverse of matrix  $P$ , use the formula for the inverse of a  $2 \times 2$  matrix:

$$P^{-1} = \frac{1}{\det(P)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

For matrix  $P = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$ , we have  $a = 5$ ,  $b = 3$ ,  $c = -1$ , and  $d = -2$ .

First, calculate the determinant of  $P$ :

$$\det(P) = (5)(-2) - (3)(-1) = -10 + 3 = -7$$

Now, find the inverse:

$$P^{-1} = \frac{1}{-7} \begin{bmatrix} -2 & -3 \\ 1 & 5 \end{bmatrix}$$

$$P^{-1} = \begin{bmatrix} \frac{2}{7} & \frac{3}{7} \\ -\frac{1}{7} & -\frac{5}{7} \end{bmatrix}$$

Thus, the inverse of  $P$  is:

$$P^{-1} = \begin{bmatrix} \frac{2}{7} & \frac{3}{7} \\ -\frac{1}{7} & -\frac{5}{7} \end{bmatrix}$$

**Hence Option 1 is the correct answer.**

---