

**TEST PAPER OF JEE(MAIN) EXAMINATION – 2019**  
**(Held On Friday 11<sup>th</sup> JANUARY, 2019) TIME : 2 : 30 PM To 5 : 30 PM**  
**MATHEMATICS**

1. If the point  $(2, \alpha, \beta)$  lies on the plane which passes through the points  $(3, 4, 2)$  and  $(7, 0, 6)$  and is perpendicular to the plane  $2x - 5y = 15$ , then  $2\alpha - 3\beta$  is equal to :-  
 (1) 5                      (2) 17                      (3) 12                      (4) 7

**Ans. (4)**

**Sol.** Normal vector of plane

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -5 & 0 \\ 4 & -4 & 4 \end{vmatrix} = -4(5\hat{i} + 2\hat{j} - 3\hat{k})$$

equation of plane is  $5(x-7) + 2y - 3(z-6) = 0$   
 $5x + 2y - 3z = 17$

2. Let  $\alpha$  and  $\beta$  be the roots of the quadratic equation  $x^2 \sin \theta - x(\sin \theta \cos \theta + 1) + \cos \theta = 0$

$(0 < \theta < 45^\circ)$ , and  $\alpha < \beta$ . Then  $\sum_{n=0}^{\infty} \left( \alpha^n + \frac{(-1)^n}{\beta^n} \right)$

is equal to :-

- (1)  $\frac{1}{1-\cos\theta} + \frac{1}{1+\sin\theta}$   
 (2)  $\frac{1}{1+\cos\theta} + \frac{1}{1-\sin\theta}$   
 (3)  $\frac{1}{1-\cos\theta} - \frac{1}{1+\sin\theta}$   
 (4)  $\frac{1}{1+\cos\theta} - \frac{1}{1-\sin\theta}$

**Ans. (1)**

**Sol.**  $D = (1 + \sin\theta \cos\theta)^2 - 4\sin\theta \cos\theta = (1 - \sin\theta \cos\theta)^2$   
 $\Rightarrow$  roots are  $\beta = \operatorname{cosec}\theta$  and  $\alpha = \cos\theta$

$$\Rightarrow \sum_{n=0}^{\infty} \left( \alpha^n + \left( -\frac{1}{\beta} \right)^n \right) = \sum_{n=0}^{\infty} (\cos\theta)^n + \sum_{n=0}^{\infty} (-\sin\theta)^n$$

$$= \frac{1}{1-\cos\theta} + \frac{1}{1+\sin\theta}$$

3. Let K be the set of all real values of x where the function  $f(x) = \sin|x| - |x| + 2(x - \pi) \cos|x|$  is not differentiable. Then the set K is equal to :-

- (1)  $\{\pi\}$                       (2)  $\{0\}$   
 (3)  $\phi$  (an empty set)      (4)  $\{0, \pi\}$

**Ans. (3)**

**Sol.**  $f(x) = \sin|x| - |x| + 2(x - \pi) \cos x$   
 $\therefore \sin|x| - |x|$  is differentiable function at  $x=0$   
 $\therefore k = \phi$

4. Let the length of the latus rectum of an ellipse with its major axis along x-axis and centre at the origin, be 8. If the distance between the foci of this ellipse is equal to the length of its minor axis, then which one of the following points lies on it ?

- (1)  $(4\sqrt{3}, 2\sqrt{3})$                       (2)  $(4\sqrt{3}, 2\sqrt{2})$   
 (3)  $(4\sqrt{2}, 2\sqrt{2})$                       (4)  $(4\sqrt{2}, 2\sqrt{3})$

**Ans. (2)**

**Sol.**  $\frac{2b^2}{a} = 8$  and  $2ae = 2b$

$$\Rightarrow \frac{b}{a} = e \text{ and } 1 - e^2 = e^2 \Rightarrow e = \frac{1}{\sqrt{2}}$$

$$\Rightarrow b = 4\sqrt{2} \text{ and } a = 8$$

so equation of ellipse is  $\frac{x^2}{64} + \frac{y^2}{32} = 1$

5. If the area of the triangle whose one vertex is at the vertex of the parabola,  $y^2 + 4(x - a^2) = 0$  and the other two vertices are the points of intersection of the parabola and y-axis, is 250 sq. units, then a value of 'a' is :-

- (1)  $5\sqrt{5}$       (2)  $(10)^{2/3}$       (3)  $5(2^{1/3})$       (4) 5

**Ans. (4)**

**Sol.** Vertex is  $(a^2, 0)$

$$y^2 = -(x - a^2) \text{ and } x = 0 \Rightarrow (0, \pm 2a)$$

$$\text{Area of triangle is } = \frac{1}{2} \cdot 4a \cdot (a^2) = 250$$

$$\Rightarrow a^3 = 125 \text{ or } a = 5$$

6. The integral  $\int_{\pi/6}^{\pi/4} \frac{dx}{\sin 2x (\tan^5 x + \cot^5 x)}$  equals :-

(1)  $\frac{1}{10} \left( \frac{\pi}{4} - \tan^{-1} \left( \frac{1}{9\sqrt{3}} \right) \right)$

(2)  $\frac{1}{5} \left( \frac{\pi}{4} - \tan^{-1} \left( \frac{1}{3\sqrt{3}} \right) \right)$

(3)  $\frac{\pi}{10}$

(4)  $\frac{1}{20} \tan^{-1} \left( \frac{1}{9\sqrt{3}} \right)$

**Ans. (1)**

**Sol.**  $I = \int_{\pi/6}^{\pi/4} \frac{dx}{\sin 2x (\tan^5 x + \cot^5 x)}$

$I = \frac{1}{2} \int_{\pi/6}^{\pi/4} \frac{\tan^4 x \sec^2 x dx}{(1 + \tan^{10} x)}$  Put  $\tan^5 x = t$

$I = \frac{1}{10} \int_{\left(\frac{1}{\sqrt{3}}\right)^5}^1 \frac{dt}{1+t^2} = \frac{1}{10} \left( \frac{\pi}{4} - \tan^{-1} \frac{1}{9\sqrt{3}} \right)$

7. Let  $(x+10)^{50} + (x-10)^{50} = a_0 + a_1x + a_2x^2 + \dots$

+  $a_{50}x^{50}$ , for all  $x \in \mathbb{R}$ , then  $\frac{a_2}{a_0}$  is equal to:-

(1) 12.50 (2) 12.00 (3) 12.75 (4) 12.25

**Ans. (4)**

**Sol.**  $(10+x)^{50} + (10-x)^{50}$

$\Rightarrow a_2 = 2 \cdot {}^{50}C_2 10^{48}, a_0 = 2 \cdot 10^{50}$

$\frac{a_2}{a_0} = \frac{{}^{50}C_2}{10^2} = 12.25$


8. Let a function  $f : (0, \infty) \rightarrow (0, \infty)$  be defined by

$f(x) = \left| 1 - \frac{1}{x} \right|$ . Then  $f$  is :-

- (1) Injective only  
(2) Not injective but it is surjective  
(3) Both injective as well as surjective  
(4) Neither injective nor surjective

**Ans. (Bonus)**

**Sol.**  $f(x) = \left| 1 - \frac{1}{x} \right| = \frac{|x-1|}{x} = \begin{cases} \frac{1-x}{x} & 0 < x \leq 1 \\ \frac{x-1}{x} & x \geq 1 \end{cases}$



$\Rightarrow f(x)$  is not injective

but range of function is  $[0, \infty)$

**Remark :** If co-domain is  $[0, \infty)$ , then  $f(x)$  will be surjective

9. Let  $S = \{1, 2, \dots, 20\}$ . A subset  $B$  of  $S$  is said to be "nice", if the sum of the elements of  $B$  is 203. Then the probability that a randomly chosen subset of  $S$  is "nice" is :-

- (1)  $\frac{6}{2^{20}}$  (2)  $\frac{5}{2^{20}}$  (3)  $\frac{4}{2^{20}}$  (4)  $\frac{7}{2^{20}}$

**Ans. (2)**

**Sol.** 7,

1,6

2,5

3,4

1,2,4

$P = \frac{5}{2^{20}}$

10. Two lines  $\frac{x-3}{1} = \frac{y+1}{3} = \frac{z-6}{-1}$  and

$\frac{x+5}{7} = \frac{y-2}{-6} = \frac{z-3}{4}$  intersect at the point  $R$ . The

reflection of  $R$  in the  $xy$ -plane has coordinates :-

- (1) (2, 4, 7) (2) (-2, 4, 7)  
(3) (2, -4, -7) (4) (2, -4, 7)

**Ans. (3)**

**Sol.** Point on  $L_1$   $(\lambda + 3, 3\lambda - 1, -\lambda + 6)$

Point on  $L_2$   $(7\mu - 5, -6\mu + 2, 4\mu + 3)$

$\Rightarrow \lambda + 3 = 7\mu - 5 \dots(i)$

$3\lambda - 1 = -6\mu + 2 \dots(ii) \Rightarrow \lambda = -1, \mu = 1$

point  $R(2, -4, 7)$

Reflection is  $(2, -4, -7)$

- 11.** The number of functions  $f$  from  $\{1, 2, 3, \dots, 20\}$  onto  $\{1, 2, 3, \dots, 20\}$  such that  $f(k)$  is a multiple of 3, whenever  $k$  is a multiple of 4, is :-

- (1)  $(15)! \times 6!$                       (2)  $5^6 \times 15$   
(3)  $5! \times 6!$                       (4)  $6^5 \times (15)!$

**Ans. (1)**

**Sol.**  $f(k) = 3m$  (3,6,9,12,15,18)  
for  $k = 4, 8, 12, 16, 20$       6.5.4.3.2 ways  
For rest numbers  $15!$  ways  
Total ways =  $6!(15)!$

- 12.** Contrapositive of the statement "If two numbers are not equal, then their squares are not equal." is :-

- (1) If the squares of two numbers are equal, then the numbers are equal.  
(2) If the squares of two numbers are equal, then the numbers are not equal.  
(3) If the squares of two numbers are not equal, then the numbers are equal.  
(4) If the squares of two numbers are not equal, then the numbers are not equal.

**Ans. (1)**

**Sol.** Contrapositive of  $p \rightarrow q$  is  $\sim q \rightarrow \sim p$

- 13.** The solution of the differential equation,

$$\frac{dy}{dx} = (x-y)^2, \text{ when } y(1) = 1, \text{ is :-}$$

- (1)  $\log_e \left| \frac{2-y}{2-x} \right| = 2(y-1)$   
(2)  $\log_e \left| \frac{2-x}{2-y} \right| = x-y$   
(3)  $-\log_e \left| \frac{1+x-y}{1-x+y} \right| = x+y-2$   
(4)  $-\log_e \left| \frac{1-x+y}{1+x-y} \right| = 2(x-1)$

**Ans. (4)**

**Sol.**  $x - y = t \Rightarrow \frac{dy}{dx} = 1 - \frac{dt}{dx}$   
 $\Rightarrow 1 - \frac{dt}{dx} = t^2 \Rightarrow \int \frac{dt}{1-t^2} = \int 1 dx$   
 $\Rightarrow \frac{1}{2} \ln \left( \frac{1+t}{1-t} \right) = x + \lambda$

$$\Rightarrow \frac{1}{2} \ln \left( \frac{1+x-y}{1-x+y} \right) = x + \lambda \quad \text{given } y(1) = 1$$

$$\Rightarrow \frac{1}{2} \ln(1) = 1 + \lambda \Rightarrow \lambda = -1$$

$$\Rightarrow \ln \left( \frac{1+x-y}{1-x+y} \right) = 2(x-1)$$

$$\Rightarrow -\ln \left( \frac{1-x+y}{1+x-y} \right) = 2(x-1)$$

- 14.** Let  $A$  and  $B$  be two invertible matrices of order  $3 \times 3$ . If  $\det(ABA^T) = 8$  and  $\det(AB^{-1}) = 8$ , then  $\det(BA^{-1}B^T)$  is equal to :-

- (1) 16                      (2)  $\frac{1}{16}$                       (3)  $\frac{1}{4}$                       (4) 1

**Ans. (2)**

**Sol.**  $|A|^2 \cdot |B| = 8$  and  $\frac{|A|}{|B|} = 8 \Rightarrow |A| = 4$  and  $|B| = \frac{1}{2}$   
 $\therefore \det(BA^{-1}B^T) = \frac{1}{4} \times \frac{1}{4} = \frac{1}{16}$

- 15.** If  $\int \frac{x+1}{\sqrt{2x-1}} dx = f(x)\sqrt{2x-1} + C$ , where  $C$  is a constant of integration, then  $f(x)$  is equal to :-

- (1)  $\frac{1}{3}(x+4)$                       (2)  $\frac{1}{3}(x+1)$   
(3)  $\frac{2}{3}(x+2)$                       (4)  $\frac{2}{3}(x-4)$

**Ans. (1)**

**Sol.**  $\sqrt{2x-1} = t \Rightarrow 2x-1 = t^2 \Rightarrow 2dx = 2t \cdot dt$

$$\int \frac{x+1}{\sqrt{2x-1}} dx = \int \frac{\frac{t^2+1}{2} + 1}{t} t dt = \int \frac{t^2+3}{2} dt$$

$$= \frac{1}{2} \left( \frac{t^3}{3} + 3t \right) = \frac{t}{6} (t^2 + 9) + c$$

$$= \sqrt{2x-1} \left( \frac{2x-1+9}{6} \right) + c = \sqrt{2x-1} \left( \frac{x+4}{3} \right) + c$$

$$\Rightarrow f(x) = \frac{x+4}{3}$$

- 16.** A bag contains 30 white balls and 10 red balls. 16 balls are drawn one by one randomly from the bag with replacement. If  $X$  be the number of white balls

drawn, the  $\left( \frac{\text{mean of } X}{\text{standard deviation of } X} \right)$  is equal to :-

- (1) 4      (2)  $\frac{4\sqrt{3}}{3}$       (3)  $4\sqrt{3}$       (4)  $3\sqrt{2}$

**Ans. (3)**

**Sol.**  $p$  (probability of getting white ball) =  $\frac{30}{40}$

$$q = \frac{1}{4} \text{ and } n = 16$$

$$\text{mean} = np = 16 \cdot \frac{3}{4} = 12$$

and standard deviation

$$= \sqrt{npq} = \sqrt{16 \cdot \frac{3}{4} \cdot \frac{1}{4}} = \sqrt{3}$$

- 17.** If in a parallelogram  $ABDC$ , the coordinates of  $A$ ,  $B$  and  $C$  are respectively  $(1, 2)$ ,  $(3, 4)$  and  $(2, 5)$ , then the equation of the diagonal  $AD$  is:-

- (1)  $5x + 3y - 11 = 0$       (2)  $3x - 5y + 7 = 0$   
(3)  $3x + 5y - 13 = 0$       (4)  $5x - 3y + 1 = 0$

**Ans. (4)**

**Sol.** co-ordinates of point  $D$  are  $(4, 7)$

$$\Rightarrow \text{line } AD \text{ is } 5x - 3y + 1 = 0$$

- 18.** If a hyperbola has length of its conjugate axis equal to 5 and the distance between its foci is 13, then the eccentricity of the hyperbola is :-

- (1) 2      (2)  $\frac{13}{6}$       (3)  $\frac{13}{8}$       (4)  $\frac{13}{12}$

**Ans. (4)**

**Sol.**  $2b = 5$  and  $2ae = 13$


$$b^2 = a^2(e^2 - 1) \Rightarrow \frac{25}{4} = \frac{169}{4} - a^2$$

$$\Rightarrow a = 6 \Rightarrow e = \frac{13}{12}$$

- 19.** The area (in sq. units) in the first quadrant bounded by the parabola,  $y = x^2 + 1$ , the tangent to it at the point  $(2, 5)$  and the coordinate axes is :-

- (1)  $\frac{14}{3}$       (2)  $\frac{187}{24}$       (3)  $\frac{37}{24}$       (4)  $\frac{8}{3}$

**Ans. (3)**



**Sol.**

$$\text{Area} = \int_0^2 (x^2 + 1) dx - \frac{1}{2} \left( \frac{5}{4} \right) (5) = \frac{37}{24}$$

- 20.** Let  $\sqrt{3}\hat{i} + \hat{j}$ ,  $\hat{i} + \sqrt{3}\hat{j}$  and  $\beta\hat{i} + (1 - \beta)\hat{j}$  respectively be the position vectors of the points  $A$ ,  $B$  and  $C$  with respect to the origin  $O$ . If the distance of  $C$  from the bisector of the acute angle between  $OA$  and  $OB$  is  $\frac{3}{\sqrt{2}}$ , then the sum of all possible values of  $\beta$  is :-

- (1) 2      (2) 1      (3) 3      (4) 4

**Ans. (2)**

**Sol.** Angle bisector is  $x - y = 0$

$$\Rightarrow \frac{|\beta - (1 - \beta)|}{\sqrt{2}} = \frac{3}{\sqrt{2}}$$

$$\Rightarrow |2\beta - 1| = 3$$

$$\Rightarrow \beta = 2 \text{ or } -1$$

- 21.** If  $\begin{vmatrix} a - b - c & 2a & 2a \\ 2b & b - c - a & 2b \\ 2c & 2c & c - a - b \end{vmatrix} = (a + b + c)(x + a + b + c)^2$ ,  $x \neq 0$  and  $a + b + c \neq 0$ , then  $x$  is equal to :-

- (1)  $-(a + b + c)$       (2)  $2(a + b + c)$   
(3)  $abc$       (4)  $-2(a + b + c)$

**Ans. (4)**

$$\text{Sol. } \begin{vmatrix} a - b - c & 2a & 2a \\ 2b & b - c - a & 2b \\ 2c & 2c & c - a - b \end{vmatrix}$$

$$R_1 \rightarrow R_1 + R_2 + R_3$$

$$= \begin{vmatrix} a + b + c & a + b + c & a + b + c \\ 2b & b - c - a & 2b \\ 2c & 2c & c - a - b \end{vmatrix}$$

$$= (a+b+c) \begin{vmatrix} 1 & 0 & 0 \\ 2b & -(a+b+c) & 0 \\ 2c & 2c & c-a-b \end{vmatrix}$$

$$= (a+b+c)(a+b+c)^2$$

$$\Rightarrow x = -2(a+b+c)$$

**22.** Let  $S_n = 1 + q + q^2 + \dots + q^n$  and

$$T_n = 1 + \left(\frac{q+1}{2}\right) + \left(\frac{q+1}{2}\right)^2 + \dots + \left(\frac{q+1}{2}\right)^n$$

where  $q$  is a real number and  $q \neq 1$ .

If  $^{101}C_1 + ^{101}C_2.S_1 + \dots + ^{101}C_{101}.S_{100} = \alpha T_{100}$ ,

then  $\alpha$  is equal to :-

- (1)  $2^{100}$       (2) 200      (3)  $2^{99}$       (4) 202

**Ans. (1)**

**Sol.**  $^{101}C_1 + ^{101}C_2.S_1 + \dots + ^{101}C_{101}.S_{100}$

$$= \alpha T_{100}$$

$$^{101}C_1 + ^{101}C_2(1+q) + ^{101}C_3(1+q+q^2) + \dots + ^{101}C_{101}(1+q+\dots+q^{100})$$

$$= 2\alpha \frac{\left(1 - \left(\frac{1+q}{2}\right)^{101}\right)}{(1-q)}$$

$$\Rightarrow ^{101}C_1(1-q) + ^{101}C_2(1-q^2) + \dots + ^{101}C_{101}(1-q^{101})$$

$$= 2\alpha \left(1 - \left(\frac{1+q}{2}\right)^{101}\right)$$

$$\Rightarrow (2^{101} - 1) - ((1+q)^{101} - 1)$$

$$= 2\alpha \left(1 - \left(\frac{1+q}{2}\right)^{101}\right)$$

$$\Rightarrow 2^{101} \left(1 - \left(\frac{1+q}{2}\right)^{101}\right) = 2\alpha \left(1 - \left(\frac{1+q}{2}\right)^{101}\right)$$

$$\Rightarrow \alpha = 2^{100}$$

**23.** A circle cuts a chord of length  $4a$  on the  $x$ -axis and passes through a point on the  $y$ -axis, distant  $2b$  from the origin. Then the locus of the centre of this circle, is :-

- (1) A hyperbola      (2) A parabola  
(3) A straight line      (4) An ellipse

**Ans. (2)**

**Sol.** Let equation of circle is

$$x^2 + y^2 + 2fx + 2fy + e = 0, \text{ it passes through } (0, 2b)$$

$$\Rightarrow 0 + 4b^2 + 2g \times 0 + 4f + c = 0$$

$$\Rightarrow 4b^2 + 4f + c = 0 \quad \dots(i)$$

$$2\sqrt{g^2 - c} = 4a \quad \dots(ii)$$

$$g^2 - c = 4a^2 \Rightarrow c = (g^2 - 4a^2)$$

Putting in equation (1)

$$\Rightarrow 4b^2 + 4f + g^2 - 4a^2 = 0$$

$$\Rightarrow x^2 + 4y + 4(b^2 - a^2) = 0, \text{ it represent a parabola.}$$

**24.** If 19th term of a non-zero A.P. is zero, then its (49th term) : (29th term) is :-

- (1) 3 : 1      (2) 4 : 1      (3) 2 : 1      (4) 1 : 3

**Ans. (1)**

**Sol.**  $a + 18d = 0 \quad \dots(1)$

$$\frac{a + 48d}{a + 28d} = \frac{-18d + 48d}{-18d + 28d} = \frac{3}{1}$$

**25.** Let  $f(x) = \frac{x}{\sqrt{a^2 + x^2}} - \frac{d-x}{\sqrt{b^2 + (d-x)^2}}$ ,  $x \in \mathbb{R}$ ,

where  $a, b$  and  $d$  are non-zero real constants.

Then :-

- (1)  $f$  is a decreasing function of  $x$   
(2)  $f$  is neither increasing nor decreasing function of  $x$   
(3)  $f'$  is not a continuous function of  $x$   
(4)  $f$  is an increasing function of  $x$

**Ans. (4)**

**Sol.**  $f(x) = \frac{x}{\sqrt{a^2 + x^2}} - \frac{d-x}{\sqrt{b^2 + (d-x)^2}}$

$$f'(x) = \frac{a^2}{(a^2 + x^2)^{3/2}} + \frac{b^2}{(b^2 + (d-x)^2)^{3/2}} > 0 \quad \forall x \in \mathbb{R}$$

$f(x)$  is an increasing function.

**26.** Let  $z$  be a complex number such that  $|z| + z = 3 + i$  (where  $i = \sqrt{-1}$ ). Then  $|z|$  is equal to :-

- (1)  $\frac{5}{4}$       (2)  $\frac{\sqrt{41}}{4}$       (3)  $\frac{\sqrt{34}}{3}$       (4)  $\frac{5}{3}$

**Ans. (4)**

**Sol.**  $|z| + z = 3 + i$

$$z = 3 - |z| + i$$

$$\text{Let } 3 - |z| = a \Rightarrow |z| = (3 - a)$$

$$\Rightarrow z = a + i \Rightarrow |z| = \sqrt{a^2 + 1}$$

$$\Rightarrow 9 + a^2 - 6a = a^2 + 1 \Rightarrow a = \frac{8}{6} = \frac{4}{3}$$

$$\Rightarrow |z| = 3 - \frac{4}{3} = \frac{5}{3}$$

- 27.** All  $x$  satisfying the inequality  $(\cot^{-1} x)^2 - 7(\cot^{-1} x) + 10 > 0$ , lie in the interval:-  
 (1)  $(-\infty, \cot 5) \cup (\cot 4, \cot 2)$   
 (2)  $(\cot 5, \cot 4)$   
 (3)  $(\cot 2, \infty)$   
 (4)  $(-\infty, \cot 5) \cup (\cot 2, \infty)$

**Ans. (4)**

**Sol.**  $\cot^{-1} x > 5, \cot^{-1} x < 2$   
 $\Rightarrow x < \cot 5, x > \cot 2$

- 28.** Given  $\frac{b+c}{11} = \frac{c+a}{12} = \frac{a+b}{13}$  for a  $\Delta ABC$  with

usual notation. If  $\frac{\cos A}{\alpha} = \frac{\cos B}{\beta} = \frac{\cos C}{\gamma}$ , then

the ordered triad  $(\alpha, \beta, \gamma)$  has a value :-

- (1) (3, 4, 5) (2) (19, 7, 25)  
 (3) (7, 19, 25) (4) (5, 12, 13)

**28. Ans. (3)**

**Sol.**  $b+c = 11\lambda, c+a = 12\lambda, a+b = 13\lambda$   
 $\Rightarrow a = 7\lambda, b = 6\lambda, c = 5\lambda$   
 (using cosine formula)

$$\cos A = \frac{1}{5}, \cos B = \frac{19}{35}, \cos C = \frac{5}{7}$$

$$\alpha : \beta : \gamma \Rightarrow 7 : 19 : 25$$

- 29.** Let  $x, y$  be positive real numbers and  $m, n$  positive integers. The maximum value of the expression

$$\frac{x^m y^n}{(1+x^{2m})(1+y^{2n})} \text{ is :-}$$

- (1)  $\frac{1}{2}$  (2)  $\frac{1}{4}$  (3)  $\frac{m+n}{6mn}$  (4) 1

**Ans. (2)**

$$\text{Sol. } \frac{x^m y^n}{(1+x^{2m})(1+y^{2n})} = \frac{1}{\left(x^m + \frac{1}{x^m}\right)\left(y^n + \frac{1}{y^n}\right)} \leq \frac{1}{4}$$

using  $AM \geq GM$

- 30.**  $\lim_{x \rightarrow 0} \frac{x \cot(4x)}{\sin^2 x \cot^2(2x)}$  is equal to :-

- (1) 2 (2) 0 (3) 4 (4) 1

$$\text{Sol. } \lim_{x \rightarrow 0} \frac{x \tan^2 2x}{\tan 4x \sin^2 x} = \lim_{x \rightarrow 0} \frac{x \left( \frac{\tan^2 2x}{4x^2} \right) 4x^2}{\left( \frac{\tan 4x}{4x} \right) 4x \left( \frac{\sin^2 x}{x^2} \right) x^2} = 1$$